


LIGAND-CONJUGATED OLIGOMERIC COMPOUNDS



FIELD OF THE INVENTION

The present invention relates to ligand-conjugated oligomeric compounds which bind to protein molecules and possess enhanced pharmacokinetic properties. The present invention further relates to methods for increasing the concentration of oligomeric compounds in serum and methods for promoting the cellular uptake of oligomeric compounds in cells.

10 BACKGROUND OF THE INVENTION

Protein synthesis is directed by nucleic acids through the intermediacy of messenger RNA (mRNA). Antisense methodology is the complementary hybridization of relatively short oligonucleotides to mRNA or DNA such that the normal, essential functions, such as protein synthesis, of these intracellular nucleic acids are disrupted. Hybridization is the sequence-specific hydrogen bonding via Watson-Crick base pairs of oligonucleotides to RNA or single-stranded DNA. Such base pairs are said to be complementary to one another.

20 The naturally-occurring events that provide the disruption of the nucleic acid function, discussed by Cohen (*Oligonucleotides: Antisense Inhibitors of Gene Expression*, CRC Press, Inc., 1989, Boca Raton, Fl.) are thought to be of

two types. The first, hybridization arrest, describes the terminating event in which the oligonucleotide inhibitor binds to the target nucleic acid and thus prevents, by simple steric hindrance, the binding of essential proteins, most often ribosomes, to the nucleic acid. Methyl phosphonate oligonucleotides (Miller *et al.* (1987) *Anti-Cancer Drug Design*, 2:117-128), and α -anomer oligonucleotides are the two most extensively studied antisense agents which are thought to disrupt nucleic acid function by hybridization arrest.

Another means by which antisense oligonucleotides disrupt nucleic acid function is by hybridization to a target mRNA, followed by enzymatic cleavage of the targeted RNA by intracellular RNase H. A 2'-deoxyribofuranosyl oligonucleotide or oligonucleotide analog hybridizes with the targeted RNA and this duplex activates the RNase H enzyme to cleave the RNA strand, thus destroying the normal function of the RNA. Phosphorothioate oligonucleotides are the most prominent example of an antisense agent that operates by this type of antisense terminating event.

Considerable research is being directed to the application of oligonucleotides and oligonucleotide analogs as antisense agents for diagnostics, research applications and potential therapeutic purposes. One of the major hurdles that has only partially been overcome *in vivo* is efficient cellular uptake which is severely hampered by the rapid degradation and excretion of oligonucleotides. The generally accepted process of cellular uptake is by receptor-mediated endocytosis which is dependent on the temperature and concentration of the oligonucleotides in serum and extra vascular fluids.

Efforts aimed at improving the transmembrane delivery of nucleic acids and oligonucleotides have utilized protein carriers, antibody carriers, liposomal delivery systems,

electroporation, direct injection, cell fusion, viral vectors, and calcium phosphate-mediated transformation. However, many of these techniques are limited by the types of cells in which transmembrane transport is enabled and by the conditions needed for achieving such transport. An alternative that is particularly attractive for the transmembrane delivery of oligonucleotides is modification of the physicochemical properties of oligonucleotides via conjugation to a molecule that facilitates transport. Another alternative is to increase the stability of oligonucleotides in serum, thereby increasing their concentration and distribution.

It has been previously reported that oligonucleotides modified with a 4-[(N-2-chloroethyl-N-methyl)amino]benzylamine reactive functionality at a 5'-phosphate position react with albumin and immunoglobulins M and G (Yu et al., *FEBS Letters*, **1994**, 334:96-98). Binding to albumin was weak at about 20 μ M with immunoglobulin binding stronger at about 4 to 6 μ M. This study further reported that oligonucleotides conjugated to steroids had increased affinity for blood cells and thus changed their distribution and increased their lifetime in serum. One method for increasing membrane or cellular transport of oligonucleotides is the attachment of a pendant lipophilic group. Ramirez et al. (*J. Am. Chem. Soc.*, **1982**, 104:5483) introduced the phospholipid group 5'-O-(1,2-di-O-myristoyl-sn-glycero-3-phosphoryl) into the dimer TpT independently at the 3' and 5' positions. Subsequently Shea et al. (*Nuc. Acids Res.*, **1990**, 18:3777) disclosed oligonucleotides having a 1,2-di-O-hexyldecyl-rac-glycerol group linked to a 5'-phosphate on the 5'-terminus of the oligonucleotide. Certain of the Shea et al. authors also disclosed these and other compounds in patent application PCT/US90/01002. A

further glucosyl phospholipid was disclosed by Guerra et al., *Tetrahedron Letters*, **1987**, 28:3581.

In other work, a cholesteryl group was attached to the internucleotide linkage between the first and second
5 nucleotides (from the 3' terminus) of an oligonucleotide. This work is disclosed in United States patent 4,958,013 and further in Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, 86:6553. Additional approaches to the delivery and study of oligonucleotides have involved the conjugation
10 of a variety of other molecules and reporter groups. The aromatic intercalating agent anthraquinone was attached to the 2' position of a sugar fragment of an oligonucleotide as reported by Yamana et al. (*Bioconjugate Chem.*, **1990**, 1:319), Lemairte et al. (*Proc. Natl. Acad. Sci. USA*, **1986**, 84:648)
15 and Leonetti et al. (*Bioconjugate Chem.*, **1990**, 1:149).

Lysine and polylysines have also been conjugated to oligonucleotides to improve their charge-size characteristics. The poly(L-lysine) was linked to the oligonucleotide via periodate oxidation of the 3'-terminal
20 ribose followed by reduction and coupling through a N-morpholine ring. Oligonucleotide-poly(L-lysine) conjugates are described in European Patent application 87109348.0. In this instance, the lysine residue was coupled to a 5' or 3' phosphate of the 5' or 3' terminal nucleotide of the
25 oligonucleotide. A disulfide linkage has also been utilized at the 3' terminus of an oligonucleotide to link a peptide to the oligonucleotide. See, Corey and Schultz, *Science*, **1987**, 238:1401; Zuckermann et al., *J. Am. Chem. Soc.*, **1988**, 110:1614; and Corey et al., *J. Am. Chem. Soc.*, **1989**,
30 111:8524.

A linking reagent for attaching biotin to the 3'-terminus of an oligonucleotide has also been described. Nelson et al., *Nuc. Acids Res.*, **1989**, 17:7187. This

reagent, N-Fmoc-O-DMT-3-amino-1,2-propanediol is now commercially available from Clontech Laboratories (Palo Alto, CA) under the name 3'-Amine on. It is also commercially available under the name 3'-Amino-Modifier

5 reagent from Glen Research Corporation (Sterling, VA). This reagent was also utilized to link a peptide to an oligonucleotide as reported by Judy et al. (*Tetrahedron Letters*, **1991**, 32:879). A similar commercial reagent (actually a series of such linkers having various lengths of

10 polymethylene connectors) for linking to the 5'-terminus of an oligonucleotide is 5'-Amino-Modifier C6. These reagents are available from Glen Research Corporation (Sterling, VA). These compounds or similar ones were utilized by Krieg et al. (*Antisense Research and Development*, **1991**, 1:161) to

15 link fluorescein to the 5'-terminus of an oligonucleotide. Other compounds of interest have also been linked to the 3'-terminus of an oligonucleotide. Asseline et al. (*Proc. Natl. Acad. Sci. USA*, **1984**, 81:3297) describe linking acridine on the 3'-terminal phosphate group of an poly (Tp)

20 oligonucleotide via a polymethylene linkage. Haralambidis et al. (*Tetrahedron Letters*, **1987**, 28:5199) report building a peptide on a solid state support and then linking an oligonucleotide to that peptide via the 3' hydroxyl group of the 3' terminal nucleotide of the oligonucleotide. Chollet

25 (*Nucleosides & Nucleotides*, **1990**, 9:957) attached an Aminolink 2 (Applied Biosystems, Foster City, CA) to the 5' terminal phosphate of an oligonucleotide. The bifunctional linking group SMPB (Pierce Chemical Co., Rockford, Il) was then used to link an interleukin protein to the

30 oligonucleotide.

Conjugation of lipids, reporters, peptides and other molecules to oligonucleotides is not limited to the terminal 3' and 5'-positions. A wide variety of conjugates have also

been reported in the literature wherein attachment is performed at any one or more of the 2'-positions on the nucleotide building blocks of the oligonucleotide. Further conjugates have also been reported wherein attachment occurs
5 on the internucleotide linkage or on one of the atoms of the nucleobase of any one of the nucleotide units of the oligonucleotide. For example, an EDTA iron complex has been linked to the 5 position of a pyrimidine nucleoside as reported by Dreyer and Dervan (*Proc. Natl. Acad. Sci. USA*,
10 **1985**, 82:968). Fluorescein has been linked to an oligonucleotide in the same manner as reported by Haralambidis et al. (*Nucleic Acid Research*, **1987**, 15:4857) and biotin in the same manner as described in PCT application PCT/US/02198. Fluorescein, biotin and pyrene
15 were also linked in the same manner as reported by Telser et al. (*J. Am. Chem. Soc.*, **1989**, 111:6966). A commercial reagent, Amino-Modifier-dT, from Glen Research Corporation (Sterling, VA) can be utilized to introduce pyrimidine nucleotides bearing similar linking groups into
20 oligonucleotides.

Manoharan et al. (PCT Application WO 93/07883) have also reported the conjugation of oligonucleotides with a variety of molecules such as steroids, reporter molecules, reporter enzymes, vitamins, non-aromatic lipophilic
25 molecules, chelators, porphyrins, intercalators, peptides and proteins through the intermediacy of varied linking groups, such as 6-aminoalkoxy and 6-aminoalkylamino groups. Conjugation has been reported at the 3'-, 5'-, 2'-, internucleotide linkage and nucleobase positions of
30 oligonucleotides. Such oligonucleotide conjugates are expected to have improved physicochemical properties that facilitated their uptake and delivery into cells as demonstrated by *in vitro* experiments. The intracellular and

intranuclear delivery of nucleic acids and oligonucleotides, however, is still a challenge. Most often, penetration of heretofore reported oligonucleotide conjugates has been found to be limited. This has typically been a problem because such conjugates have generally been designed to improve the passive absorption of the oligonucleotides where the size, physicochemical properties and extracellular concentration of the conjugate play important limiting roles. This coupled with the limited extracellular stability of nucleic acids and oligonucleotides demands the development of novel conjugates that will deliver higher levels of nucleic acids and oligonucleotides into specific tissues and targeted cells.

Albumin is the most abundant protein in mammalian systems, and plays an important role in the transport and deposition of drug substances in blood. It is generally accepted that there are two major specific drug binding sites, site I and site II on human albumin. X-ray studies of crystalline human albumin (He and Carter, *Nature*, **1992**, 358:209-215) indicate that site I and site II are located within specialized cavities in subdomain IIA and IIIA, respectively.

Interaction of oligonucleotides with proteins play an important role in absorption, distribution and pharmacokinetics. In the bloodstream, the major oligonucleotide binding proteins are immunoglobulins M and G, serum albumin, and orosomucoid α -1-acid glycoprotein (AAG). The role of plasma protein binding is an important factor in oligonucleotide disposition and efficacy. If protein binding of oligonucleotides can be modulated with small molecular conjugation, it will result in more efficacious oligonucleotide drugs.

Albumin is a water-soluble protein with a molecular weight of 66,500 comprising a single chain of 585 amino

acids containing a single tryptophan (Trp-214), low (2%) glycine content, high cystine content and a large number of charged amino acids (about 100 negative charges and 100 positive charges) and has an isoelectric point of about pH 5.0. Thus, at a plasma pH of 7.4, it has a net negative charge of -15. Nonetheless, it attracts both anions and cations. It circulates at a concentration of 3.5-5 g/100 mL in blood plasma and also exists at lower concentrations in extravascular fluids. About 60% of all human serum albumin (HSA) is located in the extravascular space (Peters, *Adv. Protein Chem.*, **1985**, 37:161). As the most abundant protein in plasma, HSA plays an important role in the maintenance of blood pH and colloidal osmotic pressure and accounts for most of the thiol content of plasma (Cys-34). Binding of drugs to albumin is usually rapidly reversible. The binding (association) constants are typically in the range of 10^4 to 10^6 M⁻¹. HSA is organized in a series of three repeating domains (I, II and III) each having two subdomains. Ligands bind to HSA generally to one or both of two binding sites. Site I is associated with the ligands warfarin, phenyl butazone. This site is localized in subdomain IIA. Site II is in subdomain IIIA and binds to diazepam and ibuprofen. Other ibuprofen analogs suprofen, pranoprofen, carprofen, fenbufen and ketoprofen, which are all non-steroidal antiinflammatory agents bind to site II. Flufenamic acid and dansylsarcosine bind to site II while dansylamide bind to site I. Barbiturates such as quinalbarbitone interact with site II and the antidiabetic tolbutamide binds to site I, site II and an unidentified site. (R)-Folinic acid binds to both sites. Other compounds that bind to HSA include thiadiazides, diazepines, and antibacterials (e.g., nalidixic acid).

Lipoproteins can contribute to the plasma binding of lipophilic drugs and dissolve in lipid core of the

lipoproteins. Cholesterol conjugated oligonucleotides are known to bind to serum proteins. Agrawal et al., ("Effect of aspirin on protein binding and tissue disposition of oligonucleotide phosphorothioate in rats," *Journal of Drug*
5 *Targeting*, **1998**, 5:303-313) describe the effect of co-administration of aspirin at a concentration of 2 mg/mL and demonstrate that the P=S oligonucleotide binding to serum albumin is reduced (as measured by % protein bound of P=S oligonucleotide). This result indicates that presence of
10 aspirin in the body or similar small molecule drugs could effectively alter protein binding of P=S oligonucleotides *in vivo*.

Pharmacokinetic studies of P=S oligonucleotide (GEM-91, 25-mer phosphorothioate oligonucleotide) in rats were
15 determined after bolus injection. One hour before administration of the drug, aspirin is administered by gavage. When P=S oligonucleotide was administered following aspirin administration in rats the following the plasma pharmacokinetic parameters ($t_{1/2} \alpha$, $t_{1/2} \beta$, AUC, etc.) were
20 lower. The tissue disposition was significantly different in that the majority of tissues. e.g. kidney, liver, spleen, bone marrow, skin, thyroid, adrenal, heart, lung, and pancreas, had lower concentrations, and gastrointestinal tissues and contents had a higher concentration. In certain
25 tissues, e.g. liver and bone marrow, the concentration of P=S oligonucleotide which was administered following aspirin administration was about half of that observed following administration of P=S oligonucleotide alone. It was seen that the rate of elimination was affected in animals
30 compared to rats receiving P=S oligonucleotide alone. A higher concentration of excreted oligonucleotide in feces from rats receiving P=S oligonucleotide following aspirin was observed compared to rats receiving P=S oligonucleotide alone. However, the effect of attaching small molecule

drugs to the oligonucleotide to modulate serum albumin binding has not been studied.

Therefore, there is a clear need for oligonucleotide conjugates having improved distribution and cellular uptake
5 and methods for their preparation, that address the shortcomings of oligonucleotide conjugates as described above. The present invention is directed to this very important end.

SUMMARY OF THE INVENTION

10 The present invention provides ligand conjugated oligomeric compounds that are capable of interacting with a protein. In particular, the ligand conjugated oligomeric compounds of the present invention bind to proteins. More particularly, the present invention provides oligomeric
15 compounds that are conjugated to drug moieties.

The oligomeric compounds of the present invention bind to serum, vascular and cellular proteins. It is preferred that the serum proteins include albumin, an immunoglobulin, a lipoprotein, α -2-macroglobulin and α -1-glycoprotein.

20 The present invention also provides ligand conjugated oligomeric compounds wherein the oligomeric compound is an oligonucleotide comprising a plurality of nucleosides. Also provided are oligonucleotides wherein the nucleosides are connected by phosphodiester linkages. Further,
25 oligonucleotides wherein the nucleosides are connected by phosphorothioate linkages are also provided. It is preferred that at least one of the nucleosides of the oligonucleotides of the present invention bear a 2'-substituent group.

30 The present invention also provides methods for increasing the concentration of an oligonucleotide in serum comprising the steps of:

(a) selecting a drug moiety that is known to bind to a

serum protein;

(b) conjugating said drug moiety to said oligonucleotide to form a conjugated oligonucleotide; and

(c) adding said conjugated oligonucleotide to said
5 serum.

The present invention further provides methods for increasing the capacity of serum for an oligonucleotide comprising the steps of:

(a) selecting a drug moiety that is known to bind to a
10 serum protein;

(b) conjugating said drug moiety to said oligonucleotide to form a conjugated oligonucleotide; and

(c) adding said conjugated oligonucleotide to said serum.

15 In one embodiment of the present invention the serum protein is a protein having a binding site for the drug moiety. In another embodiment the serum protein is a protein having a binding site for the oligonucleotide. In yet another embodiment the serum protein is a protein having
20 a binding site for the oligonucleotide and a binding site for the drug moiety such that the binding site for the oligonucleotide is distinct from the binding site for the drug moiety.

The present invention further provides methods for
25 increasing the binding of an oligonucleotide to a portion of the vascular system comprising the steps of:

(a) selecting a drug moiety that is known to bind to a protein that resides, in part, in the circulating serum and, in part, in a non-circulating portion of the vascular
30 system;

(b) conjugating said drug moiety to said oligonucleotide to form a conjugated oligonucleotide; and

(c) adding said conjugated oligonucleotide to said vascular system.

The present invention also provides methods for promoting cellular uptake of an oligonucleotide in a cell comprising the steps of:

(a) selecting a protein that resides on the cellular membrane and extends, at least in part, on the external side of said membrane;

(b) selecting a drug moiety that is known to bind to said protein;

(c) conjugating said drug moiety to said oligonucleotide to form a conjugated oligonucleotide; and

(d) exposing said cell to said conjugated oligonucleotide.

Preferably, the protein residing on the cellular membrane is a cell surface integrin.

In one embodiment of the present invention the serum protein is albumin, an immunoglobulin, α -2-macroglobulin, α -1-glycoprotein or a lipoprotein. Preferably, the serum protein is albumin.

In yet another embodiment of the present invention the drug moiety is aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepam, indomethacin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Preferably, the drug moiety is aspirin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, palmityl or carprofen. More preferably, the drug moiety is ibuprofen.

30 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph showing a comparison of HSA binding (Sigma A3782 lot 94H9318) for ibuprofen conjugates (diamonds) to unconjugated controls (triangles). Binding

curve for the phosphorothioate DNA analogs of each sequence are also shown (circles). Oligonucleotide (50 nM) was incubated with increasing concentrations of HSA as described in the text.

- 5 Figure 2 is a graph showing a comparison of the capacity of HSA (Sigma A3782 lot 97H7604) for an ibuprofen conjugate (diamonds) compared to that of an unconjugated phosphorothioate DNA (triangles). Capacity was measured at 50 mM HSA with increasing concentrations of oligonucleotide.

10 **DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS**

 The present invention provides methods of improving the pharmacokinetic properties of oligonucleotides. The invention further provides ligand conjugated oligomeric compounds having improved pharmacokinetic properties and
15 methods for their preparation. Such oligomeric compounds are prepared having covalently attached ligands that bind reversibly to one or more serum, vascular or cellular proteins. This reversible binding is expected to decrease urinary excretion, increase serum half life and greatly
20 increase the distribution of oligomeric compounds thus conjugated. The binding of particular drugs to plasma protein has been previously shown to enhance the disposition and efficacy of drugs (Herve *et al.*, *Clin. Pharmacokinet.*, **1994**, 26:44).

- 25 The therapeutic effect of an antisense oligonucleotide is realized when it interacts with a specific cellular nucleic acid and effectively negates its function. A preferred target is mRNA encoding a protein that is responsible for a disease state. To reach a target nucleic
30 acid after administration, an antisense agent should be able to overcome inherent factors such as rapid degradation in serum, short half life in serum and rapid filtration by the kidneys with subsequent excretion in the urine.

Oligonucleotides that overcome these inherent factors have increased serum half lives, distribution, cellular uptake and hence improved efficacy. These enhanced pharmacokinetic parameters have been shown for selected drug molecules that
5 bind plasma proteins (Olson and Christ, *Annual Reports in Medicinal Chemistry*, **1996**, 31:327). Two proteins that have been studied more than most are human serum albumin (HSA) and α -1-acid glycoprotein. HSA binds a variety of endogenous and exogenous ligands with association constants
10 typically in the range of 10^4 to 10^6 M⁻¹. Association constants for ligands with α -1-acid glycoprotein are similar to those for HSA.

At least for therapeutic purposes, antisense oligonucleotides should have a degree of stability in serum
15 to allow distribution and cellular uptake. The prolonged maintenance of therapeutic levels of antisense agents in serum will have a significant effect on the distribution and cellular uptake and unlike conjugate groups that target specific cell receptors the increased serum stability will
20 effect all cells. Numerous efforts have focused on increasing the cellular uptake of oligonucleotides including increasing the membrane permeability via conjugates and cellular delivery of oligonucleotides.

Many drugs reversibly bind to plasma proteins. A
25 representative list, which is not meant to be inclusive, includes: aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, benzothiadiazides,
30 chlorothiazide, diazepines (such as for example fludiazepam and diazepam) indomethacin, barbiturates (such as for example quinalbarbitone), cephalosporins, sulfa drugs, antidiabetics (such as for example tolbutamide), antibacterials (such as for example a group of quinolones;

nalidixic acid and cinoxacin) and several antibiotics. Serum albumin is the most important protein among all plasma proteins for drug binding, although binding to other proteins (for example, macroglobulin G₂, immunoglobulins, lipoproteins, alpha-1-acid glycoprotein, thrombin) is also important.

Ligands that bind serum, vascular or cellular proteins may be attached via an optional linking moiety to one or more sites on an oligonucleotide of the invention. These sites include one or more of, but are not limited to, the 2'-position, 3'-position, 5'-position, the internucleotide linkage, and a nucleobase atom of any nucleotide residue. The attachment of ligands to such structures can be performed, according to some preferred embodiments of the invention, using a linking group, or without the use of such a linking group.

In some preferred embodiments of the invention, one or more protein binding ligands are attached to an oligonucleotide via linking groups, to form a ligand conjugated oligonucleotide. Preferred linking groups of the invention include, but are not limited to, 6-aminoalkoxy linkers, 6-aminoalkylamino linkers, cysteamine, heterobifunctional linkers, homobifunctional linkers, and a universal linker (derived from 3-dimethoxytrityloxy-2-aminopropanol). A particularly preferred linking group for the synthesis of ligand conjugated oligonucleotides of the invention is a 6-aminohexyloxy group. A variety of heterobifunctional and homobifunctional linking moieties are available from Pierce Co. (Rockford, IL). Such heterobifunctional and homobifunctional linking moieties are particularly useful in conjunction with the 6-aminoalkoxy and 6-aminoalkylamino moieties to form extended linkers useful for linking ligands to a nucleoside. Further useful linking groups that are commercially available are 5'-Amino-

Modifier C6 and 3'-Amino-Modifier reagents, both available from Glen Research Corporation (Sterling, VA). 5'-Amino-Modifier C6 is also available from ABI (Applied Biosystems Inc., Foster City, CA) as Aminolink-2, while the 3'-Amino-
5 Modifier is also available from Clontech Laboratories Inc. (Palo Alto, CA). In addition, a nucleotide analog bearing a linking group pre-attached to the nucleoside is commercially available from Glen Research Corporation under the tradename "Amino-Modifier-dT." This nucleoside-linking group reagent,
10 a uridine derivative having an [N(7-trifluoroacetyl-amino-heptyl)3-acrylamido] substituent group at the 5 position of the pyrimidine ring, is synthesized as per the procedure of Jablonski *et al.* (*Nucleic Acid Research*, 1986, 14:6115). The present invention also includes as nucleoside analogs
15 adenine nucleosides functionalized to include a linker on the N6 purine amino group, guanine nucleosides functionalized to include a linker at the exocyclic N2 purine amino group, and cytosine nucleosides functionalized to include a linker on either the N4 pyrimidine amino group
20 or the 5 pyrimidine position. Such nucleoside analogs are incorporated into oligonucleotides with a ligand attached to the linker either pre- or post-oligomerization.

In a preferred embodiment of the present invention ligand molecules are selected for conjugation to
25 oligonucleotides on the basis of their affinity for one or more proteins. These proteins may be serum, vascular or cellular proteins. Serum proteins are proteins that are present in the fluid portion of the blood, obtained after coagulation and removal of the fibrin clot and blood cells,
30 as distinguished from the plasma in circulating blood. Vascular proteins are proteins that are present in portions of the vascular system relating to or containing blood vessels. Cellular proteins are membrane proteins which have at least a portion of the protein extending extracellularly


and assisting in the process of endocytosis.

Many ligands having an affinity for proteins are well documented in the literature and are amenable to the present invention. A preferred group of ligands are small molecules including drug moieties. According to the present invention, drug moieties include, but are not limited to, warfarin and coumarins including substituted coumarins, isocoumarin derivatives, 7-anilinocoumarin-4-acetic acid, profens including ibuprofen, enantiomers of ibuprofen (r-ibuprofen and s,-ibuprofen), ibuprofen analogs, ketoprofen, carprofen, etodolac, suprofen, indoprofen, fenbufen, arylpropionic acids, arylalkanoic acids, 2-aryl-2-fluoropropionic acids, glibenclamide, acetohexamide, arylalkanoic acids, tolbutamide, gliclazide, metformin, curcumin, digitoxin, digoxin, diazepam, benzothiadiazides, chlorothiazide, diazepines, benzodiazepines, naproxen, phenyl butazone, oxyphenbutazone, dansyl amide, dansylsarcosine, 2,3,5-triiodobenzoic acid, palmitic acid, aspirin, salicylates, substituted salicylates, penicillin, flurbiprofen, pirprofen, oxaprozin, flufenamic acid, deoxycholic acid, glycyrrhizin, azathioprine, butibufen, ibufenac, 5-fluoro-1-typtaphan, 5-fluoro-salicylic, acidazapropanazone, mefenamic acid, indomethacin, flufenamic acid, bilirubin, ibuprofen, lysine complexes, diphenyl, hydantoin, valproic acid, tolmetin, barbiturates (such as, for example, quinalbarbitone), cephalosporins, sulfa drugs, antidiabetics (such as, for example, tollbutamide), antibacterials (such as, for example, quinolones, nalidixic acid and cinoxacin) and several antibiotics.

30 In one embodiment of the present invention the drug moiety bears a carboxylic acid group. In another embodiment of the present invention the drug moiety is a propionic acid derivative.

In another preferred embodiment, the drug moiety is an

arylpropionic acid of the formula:



wherein:

one of R₁ and R₂ is C₁ to C₁₂ alkyl and the other of R₁ and
5 R₂ is aryl; or

both R₁ and R₂ are C₁ to C₁₂ alkyl; or

both R₁ and R₂ are aryl.

In preferred embodiments, the aryl group may be substituted or unsubstituted benzyl, phenyl, xylyl, naphthyl,
10 toluyl, pyrenyl, anthracyl, phenanthryl, azulyl, phenethyl, cinnamyl, benzhydryl, and mesityl wherein said substituents are hydroxyl, alkyl, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, or alkyl, substituted alkyl, aryl, alkenyl, or alkynyl groups.

15 In one preferred embodiment of the invention the protein targeted by a ligand conjugated oligomeric compound is a serum protein. It is preferred that the serum protein targeted by a conjugated oligomeric compound is an immunoglobulin (an antibody). Preferred immunoglobulins are immunoglobulin G and
20 immunoglobulin M. Immunoglobulins are known to appear in blood serum and tissues of vertebrate animals.

In another embodiment of the invention the serum protein targeted by a conjugated oligomeric compound is a lipoprotein. Lipoproteins are blood proteins having molecular weights
25 generally above 20,000 that carry lipids and are recognized by specific cell surface receptors. The association with lipoproteins in the serum will initially increase pharmacokinetic parameters such as half life and distribution. A secondary consideration is the ability of lipoproteins to
30 enhance cellular uptake via receptor-mediated endocytosis.

In yet another embodiment the serum protein targeted by

a ligand conjugated oligomeric compound is α -2-macroglobulin. In yet a further embodiment the serum protein targeted by a ligand conjugated oligomeric compound is α -1-glycoprotein.

As used herein, the term "protected" means that the indicated moiety has a protecting group appended thereon. In some preferred embodiments of the invention compounds contain one or more protecting groups. A wide variety of protecting groups can be employed in the methods of the invention. In general, protecting groups render chemical functionalities inert to specific reaction conditions, and can be appended to and removed from such functionalities in a molecule without substantially damaging the remainder of the molecule.

Representative hydroxyl protecting groups, for example, are disclosed by Beaucage et al. (*Tetrahedron*, **1992**, 48:2223-2311). Further hydroxyl protecting groups, as well as other representative protecting groups, are disclosed in Greene and Wuts, *Protective Groups in Organic Synthesis*, Chapter 2, 2d ed., John Wiley & Sons, New York, 1991, and *Oligonucleotides And Analogues A Practical Approach*, Ekstein, F. Ed., IRL Press, N.Y., 1991, each of which is hereby incorporated by reference in its entirety.

Examples of hydroxyl protecting groups include, but are not limited to, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p'-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, triphenylsilyl, benzoylformate, acetate, chloroacetate, trichloroacetate, trifluoroacetate, pivaloate, benzoate, p-phenylbenzoate, 9-fluorenylmethyl carbonate, mesylate and tosylate.

Amino-protecting groups stable to acid treatment are selectively removed with base treatment, and are used to make reactive amino groups selectively available for substitution.

Examples of such groups are the Fmoc (E. Atherton and R.C. Sheppard in *The Peptides*, S. Udenfriend, J. Meienhofer, Eds., Academic Press, Orlando, 1987, volume 9, p.1) and various substituted sulfonylethyl carbamates exemplified by the Nsc
5 group (Samukov et al., *Tetrahedron Lett*, **1994**, 35:7821; Verhart and Tesser, *Rec. Trav. Chim. Pays-Bas*, **1987**, 107:621).

Additional amino-protecting groups include, but are not limited to, carbamate-protecting groups, such as 2-trimethylsilylethoxycarbonyl (Teoc), 1-methyl-1-(4-
10 biphenyl)ethoxycarbonyl (Bpoc), t-butoxycarbonyl (BOC), allyloxycarbonyl (Alloc), 9-fluorenylmethyloxycarbonyl (Fmoc), and benzyloxycarbonyl (Cbz); amide-protecting groups, such as formyl, acetyl, trihaloacetyl, benzoyl, and nitrophenylacetyl; sulfonamide-protecting groups, such as 2-nitrobenzenesulfonyl;
15 and imine- and cyclic imide-protecting groups, such as phthalimido and dithiasuccinoyl. Equivalents of these amino-protecting groups are also encompassed by the compounds and methods of the present invention.

In a preferred embodiment of the present invention
20 oligonucleotides are provided including a number of linked nucleosides wherein at least one of the nucleosides is a 2'-functionalized nucleoside having a ligand molecule linked to the 2'-position of the nucleoside; a heterocyclic base functionalized nucleoside having a ligand molecule linked to
25 the heterocyclic base of the nucleoside, a 5' terminal nucleoside having a ligand molecule linked to the 5'-position of the nucleoside, a 3' terminal nucleoside having a ligand molecule linked to the 3'-position of the nucleoside, or an inter-strand nucleoside having a ligand molecule linked to an
30 inter-stand linkage linking said inter-strand nucleoside to an adjacent nucleoside.

Ligand conjugated oligonucleotides of the invention may be synthesized by the use of an oligonucleotide that bears a pendant reactive functionality such as that derived from the

attachment of a linking molecule onto the oligonucleotide. This reactive oligonucleotide may be reacted directly with commercially available ligands, ligands that are synthesized bearing a variety of protecting groups, or ligands that have
5 a linking moiety attached thereto. The methods of the present invention facilitate the synthesis of ligand conjugated oligonucleotides by the use of, in some preferred embodiments, nucleoside monomers that have been appropriately conjugated with ligands and that may further be attached to a solid
10 support material. Such ligand-nucleoside conjugates optionally attached to a solid support material are prepared according to some preferred embodiments of the methods of the present invention via reaction of a selected serum binding ligand with a linking moiety located on a 2', 3', or 5' position of a
15 nucleoside or oligonucleotide.

The present invention provides methods for increasing the concentration of an oligonucleotide in serum. According to such methods, a drug moiety that is known to bind to a serum protein is selected and conjugated to an oligonucleotide, thus
20 forming a conjugated oligonucleotide. This conjugated oligonucleotide is then added to the serum.

The present invention further provides methods for increasing the capacity of serum for an oligonucleotide. According to such methods, a drug moiety that is known to bind
25 to a serum protein is selected and conjugated to an oligonucleotide, thus forming a conjugated oligonucleotide. This conjugated oligonucleotide is then added to the serum.

The present invention also provides methods for increasing the binding of an oligonucleotide to a portion of the vascular system. According to such methods, a drug moiety that is known
30 to bind to a vascular protein is selected. The vascular protein selected is a protein which resides, in part, in the circulating serum and, in part, in the non-circulating portion of the vascular system. This drug moiety is conjugated to an

oligonucleotide to form a conjugated oligonucleotide, which is then added to the vascular system.

The present invention further provides methods for promoting the cellular uptake of an oligonucleotide in a cell. According to such methods, a cellular protein is selected. This cellular protein is a protein that resides on the cellular membrane and extends, in part, extracellularly so that part of this cellular protein extends onto the external side of the cellular membrane. Next, a drug moiety that is known to bind to the cellular protein is selected and conjugated to an oligonucleotide to form a conjugated oligonucleotide. This conjugated oligonucleotide is then brought into contact with cells in which cellular uptake of the oligonucleotide is to be promoted.

The present invention also provides methods of increasing cellular uptake of an oligonucleotide comprising contacting an organism with an oligonucleotide of the invention, said oligonucleotide being conjugated to a ligand.

Ligand conjugated oligomeric compounds of the present invention can be included in compositions that further include one or more inert carrier compounds.

Antisense therapeutics can be practiced in a plethora of various organisms ranging from unicellular prokaryotic and eukaryotic organisms to multicellular eukaryotic organisms. Any organism that utilizes DNA-RNA transcription or RNA-protein translation as a fundamental part of its hereditary, metabolic or cellular control is susceptible to antisense therapeutics and/or prophylactics. Seemingly diverse organisms such as bacteria, yeast, protozoa, algae, all plant and all higher animal forms, including warm-blooded animals, can be treated by antisense therapy. Further, since each of the cells of multicellular eukaryotes also includes both DNA-RNA transcription and RNA-protein translation as an integral part of its cellular activity, antisense therapeutics and/or

diagnostics can also be practiced on such cellular populations. Furthermore, many of the organelles, e.g. mitochondria and chloroplasts, of eukaryotic cells also include transcription and translation mechanisms. As such, single cells, cellular
5 populations or organelles can also be included within the definition of organisms that are capable of being treated with antisense therapeutics or diagnostics. As used herein, therapeutics is meant to include both the eradication of a disease state, killing of an organism, e.g. bacterial,
10 protozoan or other infection, or control of erratic or harmful cellular growth or expression.

In a preferred embodiment of the present invention a ligand having an affinity for a serum protein is attached to at least one nucleoside in an antisense diagnostic or
15 therapeutic agent to enhance the pharmacokinetic properties of the antisense therapeutic or diagnostic agent. Such improved pharmacokinetic properties include, but are not limited to, increased binding of the antisense compound to serum proteins, increased plasma concentration of the antisense compound,
20 increased tissue distribution, increased capacity of binding of the antisense compound to serum proteins and increased half-lives. Such an antisense diagnostic or therapeutic agent is preferably a nucleic acid or oligonucleotide formed of a plurality of linked nucleosides of a sequence that are
25 "antisense" to a region of an RNA or DNA of interest. The nucleosides are linked by phosphorus-containing or non-phosphorus-containing covalent internucleoside linkages. One or more nucleosides of the oligonucleotide are conjugated to include a ligand molecule bound to the nucleoside with or
30 without a linking group. For the purposes of identification, such conjugated nucleosides can be characterized as ligand bearing nucleosides or ligand-nucleoside conjugates. The linked nucleosides having at least one conjugated nucleoside within their sequence will demonstrate enhanced antisense

activity when compared to like linked nucleoside or oligonucleotides of the same sequence that are not conjugated.

The ligand conjugated oligonucleotides of the present invention also include conjugates of oligonucleotides and
5 linked nucleosides wherein the ligand is attached directly onto the nucleoside or nucleotide without the intermediacy of a linker group. This attachment of ligand may be performed at either one or more of the 2'-, 3'-, 5'-, nucleobase or internucleoside linkage positions of the oligonucleotide or
10 linked nucleosides of the invention. Ligands may preferably be attached, via linking groups, at a carboxyl, amino or oxo groups of the ligand. Typical linking groups may be ester, amide or carbamate groups.

In the context of this invention, the terms "oligomer" and
15 "oligomeric compound" refer not only to a plurality of naturally-occurring or non-naturally-occurring nucleosides joined together in a specific sequence but are further inclusive of all types of oligomeric compounds currently known in the art such as oligonucleotide analogs, peptide nucleic
20 acids and locked nucleic acids where the sugar is involved in locking the conformation. Many diverse motifs are currently known for improving the desired effect for the specific target that the oligomeric compound is aimed at such as for example chimeric oligomeric compounds where there are more than one
25 type of internucleoside linkages dividing the oligomeric compound into regions. Oligomeric compounds are typically structurally distinguishable from, yet functionally interchangeable with, naturally-occurring or synthetic wild-type oligonucleotides. Thus, oligomeric compounds include all such
30 structures which function effectively to mimic the structure and/or function of a desired RNA or DNA strand, for example, by hybridizing to a target. Whereas the term "oligonucleotide" has a well defined meaning in the art, the term "oligomeric compound" or "oligomer" is intended to be broader, inclusive

of oligomers having all manner of modifications known in the art. Gapped or chimeric compounds are disclosed in for example, U.S. Patent No. 5,623,065, issued April 22, 1997, the contents of which are incorporated herein by reference.

5 In the context of this invention, the term "oligomeric compound" includes linked nucleosides having phosphorus and non-phosphorus linkages and mixed backbone oligomers. A representative list of phosphorus containing and non-phosphorus containing linkages amenable to the present invention includes:

10 phosphorus containing linkages

phosphorodithioate (-O-P(S)(S)-O-);

phosphorothioate (-O-P(S)(O)-O-);

phosphoramidate (-O-P(O)(NJ)-O-);

phosphonate (-O-P(J)(O)-O-);

15 phosphotriesters (-O-P(O)(J)(O)-O-);

phosphoramidate (-O-P(O)(NJ)-S-);

thionoalkylphosphonate (-O-P(S)(J)-O-);

thionoalkylphosphotriester (-O-P(O)(OJ)-S-);

boranophosphate (-R⁵-P(O)(O)-J-);

20 non-phosphorus containing linkages

thiodiester (-O-C(O)-S-);

thionocarbamate (-O-C(O)(NJ)-S-);

siloxane (-O-Si(J)₂-O-);

carbamate (-O-C(O)-NH- and -NH-C(O)-O-)

25 sulfamate (-O-S(O)(O)-N- and -N-S(O)(O)-N-;

morpholino sulfamide (-O-S(O)(N(morpholino))-);

sulfonamide (-O-SO₂-NH-);

sulfide (-CH₂-S-CH₂-);

sulfonate (-O-SO₂-CH₂-);

30 N,N'-dimethylhydrazine (-CH₂-N(CH₃)-N(CH₃)-);

thioformacetal (-S-CH₂-O-);

formacetal (-O-CH₂-O-);

thioketal (-S-C(J)₂-O-); and

ketal (-O-C(J)₂-O-);

amine (-NH-CH₂-CH₂-);
hydroxylamine (-CH₂-N(J)-O-);
hydroxyimine (-CH=N-O-); and
hydrazinyl (-CH₂-N(H)-N(H)-).

5 where "J" denotes a substituent group which is commonly hydrogen or an alkyl group or a more complicated group that varies from one type of linkage to another.

10 In addition to linking groups as described above that involve the modification or substitution of the -O-P-O- atoms of a naturally occurring linkage, included within the scope of the present invention are linking groups that include modification of the 5'-methylene group as well as one or more of the -O-P-O- atoms. Linkages of this type are well documented in the prior art and include without limitation the
15 following:

amides (-CH₂-CH₂-N(H)-C(O)) and -CH₂-O-N=CH-; and
alkylphosphorus (-C(J)₂-P(=O)(OJ)-C(J)₂-C(J)₂-).
wherein J is as described above.

Synthetic schemes for the synthesis of the substitute
20 internucleoside linkages described above are disclosed in: WO 91/08213; WO 90/15065; WO 91/15500; WO 92/20822; WO 92/20823; WO 91/15500; WO 89/12060; EP 216860; PCT/US 92/04294; PCT/US 90/03138; PCT/US 91/06855; PCT/US 92/03385; PCT/US 91/03680; U.S. Application Nos. 07/990,848; 07,892,902; 07/806,710;
25 07/763,130; 07/690,786; U.S. Patent Nos. 5,466,677; 5,034,506; 5,124,047; 5,278,302; 5,321,131; 5,519,126; 4,469,863; 5,455,233; 5,214,134; 5,470,967; 5,434,257; Stirchak, E.P., et al., *Nucleic Acid Res.*, **1989**, 17, 6129-6141; Hewitt, J.M., et al., **1992**, 11, 1661-1666; Sood, A., et al., *J. Am. Chem. Soc.*,
30 **1990**, 112, 9000-9001; Vaseur, J.J. et al., *J. Amer. Chem. Soc.*, **1992**, 114, 4006-4007; Musichi, B., et al., *J. Org. Chem.*, **1990**, 55, 4231-4233; Reynolds, R.C., et al., *J. Org. Chem.*, **1992**, 57, 2983-2985; Mertes, M.P., et al., *J. Med. Chem.*, **1969**, 12, 154-

157; Mungall, W.S., et al., *J. Org. Chem.*, **1977**, 42, 703-706; Stirchak, E.P., et al., *J. Org. Chem.*, **1987**, 52, 4202-4206; Coull, J.M., et al., *Tet. Lett.*, **1987**, 28, 745; and Wang, H., et al., *Tet. Lett.*, **1991**, 32, 7385-7388.

5 Other modifications can be made to the sugar, to the base, or to the phosphate group of the nucleotide. Representative modifications are disclosed in International Publication Numbers WO 91/10671, published July 25, 1991, WO 92/02258, published February 20, 1992, WO 92/03568, published March 5,
10 1992, and United States Patents 5,138,045, 5,218,105, 5,223,618 5,359,044, 5,378,825, 5,386,023, 5,457,191, 5,459,255, 5,489,677, 5,506,351, 5,541,307, 5,543,507, 5,571,902, 5,578,718, 5,587,361, 5,587,469, all assigned to the assignee of this application. The disclosures of each of the above
15 referenced publications are herein incorporated by reference.

Another modification that is used to prepare oligomeric compounds amenable to the present invention includes LNA's (locked nucleic acid) which are novel conformationally restricted oligonucleotide analogs containing 2'-O,4'-C-
20 methylene LNA nucleoside monomers. LNA and LNA analogs display very high duplex thermal stabilities with complementary DNA and RNA ($T_m = +3$ to $+10$ C), stability towards 3'-exonucleolytic degradation, and good solubility properties (see for example:

LNA (locked nucleic acids, synthesis and high-affinity
25 nucleic acid recognition, Singh et al., *Dep. Chem., Univ. Copenhagen, Copenhagen, Den. Chem. Commun., (Cambridge)* (1998), (4), 455-456. A novel class of nucleic acid analogs, termed LNA (locked nucleic acids), is introduced. Following the Watson-Crick base pairing rules, LNA forms duplexes with
30 complementary DNA and RNA with remarkably increased thermal stabilities and generally improved selectivity.

Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclonucleoside monomers,

oligomerization, and nucleic acid recognition properties have been described (see Koshkin *et al.*, Department of Chemistry, University of Copenhagen, Copenhagen, Den. Tetrahedron (1998), 54(14), 3607-3630. LNA monomers, have been synthesized and
5 their nucleic acid recognition potential evaluated for six different nucleobases, namely adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil. Studies of mis-matched sequences show that LNA obey the Watson-Crick base pairing rules with generally improved selectivity compared to the
10 corresponding unmodified reference strands.

Potent and nontoxic antisense oligonucleotides containing locked nucleic acids have been described (Wahlestedt *et al.*, Center for Genomics Research, Karolinska Institutet, Stockholm, Swed., Proc. Natl. Acad. Sci. U. S. A. (2000), 97(10), 5633-
15 5638.) The authors have demonstrated locked nucleic acid (LNA), confers several desired properties to antisense agents. LNA/DNA copolymers were not degraded readily in blood serum and cell extracts. LNA/DNA copolymers exhibited potent antisense activity on assay systems as disparate as a G-protein-coupled
20 receptor in living rat brain and an Escherichia coli reporter gene.

The conformations of locked nucleic acids (LNA) has been determined by Petersen *et al.*, Department of Chemistry, University of Southern Denmark, Odense University, Odense,
25 Den., J. Mol. Recognit., (2000), 13(1), 44-53. The authors have used 2D NMR spectroscopy to show that the locked conformation of the LNA nucleotides both in ssLNA and in the duplexes organize the phosphate backbone in such a way as to introduce higher population of the N-type conformation. These
30 conformational changes are associated with an improved stacking of the nucleobases.

LNA properties have been described by Wengel *et al.*, Center For Synthetic Bioorganic Chemistry, Department of Chemistry, University of Copenhagen, Copenhagen, Den.

Nucleosides Nucleotides (1999), 18(6 & 7), 1365-1370.

LNA forms duplexes with complementary DNA, RNA or LNA with high thermal affinities. CD spectra show that duplexes involving fully modified LNA (esp. LNA:RNA) structurally resemble an A-form RNA:RNA duplex. NMR examination of an LNA:DNA duplex confirm the 3'-endo conformation of an LNA monomer. Recognition of double-stranded DNA is demonstrated suggesting strand invasion by LNA. Lipofectin-mediated efficient delivery of LNA into living human breast cancer cells has been accomplished.

In a patent application, preparation of locked nucleoside analogs-containing oligodeoxyribonucleotide duplexes as substrates for nucleic acid polymerases has been described

Wengel, Jesper; Nielsen, Poul. (Exiqon A/S, Den.). PCT Int. Appl. (1999), 269 pp Patent written in English. Application: WO 98-DK393 19980914. Priority: DK 97-1054 19970912; DK 97-1492 19971219; DK 98-61 19980116; DK 98-286 19980303; DK 98-585 19980429; US 98-88309 19980605. CAN 130:252609 AN 1999:216926

Bicyclic and tricyclic nucleoside and nucleotide analogs were prepared as well as oligodeoxyribonucleotides comprising such elements I (B is selected from hydrogen, hydroxy, alkoxy, alkyl, acyloxy, nucleobases, DNA intercalators; P designates the radical position for an internucleoside linkage to a succeeding monomer, or a 5'-terminal group, such internucleoside linkage or 5'-terminal group optionally including the substituent R5; X is selected from O, S, substituted N, substituted C; R1, R1*, R2, R2*, R3, R3*, R4*, R5, R5*, are biradical(s), independently selected from hydrogen, alkyl, alkenyl, alkynyl, hydroxy, alkoxy, alkenyloxy, carboxy, alkoxycarbonyl, alkylcarbonyl, formyl, aryl, aryloxy-carbonyl, aryloxy, arylcarbonyl, heteroaryl, carbamido, alkanoyloxy, sulfono, alkylsulfonyloxy, nitro, azido, sulphanyl, alkylthio, halogen, DNA intercalators). Thus,

(1S,5R,6R,8R)-5-(2-cyanoethoxy(diisopropylamino)phosphinoxy)-6-(4,4'-dimethoxytrityloxymethyl)-8-(thymine-1-yl)-2,7-dioxabicyclo[3.3.0]nonane was prepd. and incorporated into oligodeoxyribonucleotides. The nucleotide analogs, LNAs
5 (Locked Nucleoside Analogs), are able to provide valuable improvements to oligonucleotides with respect to affinity and specificity towards complementary RNA and DNA oligomers. The novel type of LNA modified oligonucleotides, as well as the LNAs as such, are useful in a wide range of diagnostic
10 applications as well as therapeutic applications. Among these can be mentioned antisense applications, PCR applications, strand displacement oligomers, as substrates for nucleic acid polymerases, as nucleotide based drugs, etc.

LNA has been shown to form exceedingly stable LNA:LNA
15 Duplexes (Koshkin et al., Center for Synthetic Bioorganic Chemistry Department of Chemistry, University of Copenhagen, Copenhagen, Den. J. Am. Chem. Soc. (1998), 120(50), 13252-13253).

LNA:LNA hybridization was shown to be the most thermally
20 stable nucleic acid type duplex system, and the RNA-mimicking character of LNA was established at the duplex level. Introduction of 3 LNA monomers (T or A) induced significantly increase melting points ($T_m = +15/+11$) toward DNA complements. The universality of LNA-mediated hybridization has been
25 stressed by the formation of exceedingly stable LNA:LNA duplexes. The RNA-mimicking of LNA was reflected with regard to the N-type conformational restriction of the monomers and to the secondary structure of the LNA:RNA duplex.

Synthesis of 2'-amino-LNA, a novel conformationally
30 restricted high-affinity oligonucleotide analog with a Handle has been shown (see Singh et al., Center for Synthetic Bioorganic Chemistry Department of Chemistry, University of Copenhagen, Copenhagen, Den. J. Org. Chem. (1998), 63(26), 10035-10039.)

2'-Amino- and 2'-methyldamino-locked nucleic acids (2'-amino-LNA) containing monomer nucleoside I ($R = \text{Me}, \text{COCF}_3$) were prepared and thermal stability of their duplexes with complementary RNA and DNA strands have been previously
5 reported.

Similarly, The first analogs of LNA, phosphorothioate-LNA and 2'-thio-LNAs have been prepared (see Kumar *et al.*, Center for Synthetic Bioorganic Chemistry, Department of Chemistry, University of Copenhagen, Copenhagen, Den. Bioorg. Med. Chem.
10 Lett. (1998), 8(16), 2219-2222.)

Synthesis of the chemically modified LNA analogs has also been reported. A 9-mer phosphorothioate-LNA containing three LNA thymine monomers ($I, X = \text{O}, Y = \text{S}, R = \text{Me}$) and 9-mer LNAs containing one, three or five 2'-thio-LNA monomers ($I, X = \text{S},$
15 $Y = \text{O}, R = \text{H}$) were able to recognize both complementary DNA and RNA with thermal affinities comparable to those of parent LNA.

Synthesis of novel bicyclo[2.2.1] ribonucleosides, 2'-amino- and 2'-thio-LNA monomeric nucleosides has been described in Singh *et al.*, Center for Synthetic Bioorganic Chemistry
20 Department of Chemistry Chemical Laboratory II, University of Copenhagen, Copenhagen, Den. J. Org. Chem. (1998), 63(18), 6078-6079.)

The present invention employs oligonucleotides for use in antisense modulation of the function of DNA or messenger RNA
25 (mRNA) encoding a protein the modulation of which is desired, and ultimately to regulate the amount of such a protein. Hybridization of an antisense oligonucleotide with its mRNA target interferes with the normal role of mRNA and causes a modulation of its function in cells. The functions of mRNA to
30 be interfered with include all vital functions such as translocation of the RNA to the site for protein translation, actual translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, turnover or degradation of the mRNA and possibly even independent catalytic activity which

may be engaged in by the RNA. The overall effect of such interference with mRNA function is modulation of the expression of a protein, wherein "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of
5 the protein. In the context of the present invention, inhibition is the preferred form of modulation of gene expression.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of
10 ribonucleic acid or deoxyribonucleic acid. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent intersugar (backbone) linkages as well as modified oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or
15 substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced binding to target and increased stability in the presence of nucleases. The oligonucleotides of the present invention preferably comprise
20 from about 5 to about 50 nucleosides. It is more preferred that such oligonucleotides comprise from about 8 to about 30 nucleosides, with 15 to 25 nucleosides being particularly preferred.

An oligonucleotide is a polymer of repeating units
25 generically known as nucleotides or nucleosides. An unmodified (naturally occurring) nucleotide has three components: (1) a nitrogenous base linked by one of its nitrogen atoms to (2) a 5-carbon cyclic sugar and (3) a phosphate, esterified to carbon 5 of the sugar. When incorporated into an oligonucleotide
30 chain, the phosphate of a first nucleotide is also esterified to carbon 3 of the sugar of a second, adjacent nucleotide. The "backbone" of an unmodified oligonucleotide consists of (2) and (3), that is, sugars linked together by phosphodiester linkages between the C5 (5') position of the sugar of a first nucleotide

and the C3 (3') position of a second, adjacent nucleotide. A "nucleoside" is the combination of (1) a nucleobase and (2) a sugar in the absence of a phosphate moiety (Kornberg, *DNA Replication*, W.H. Freeman & Co., San Francisco, 1980, pages 4-5 7). The backbone of an oligonucleotide positions a series of bases in a specific order; the written representation of this series of bases, which is conventionally written in 5' to 3' order, is known as a nucleotide sequence.

Oligonucleotides may comprise nucleoside or nucleotide sequences sufficient in identity and number to effect specific hybridization with a particular nucleic acid. Such oligonucleotides which specifically hybridize to a portion of the sense strand of a gene are commonly described as "antisense." In the context of the invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleosides or nucleotides. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that an oligonucleotide need not be 100% complementary to its target

DNA sequence to be specifically hybridizable. An oligonucleotide is specifically hybridizable when binding of the oligonucleotide to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a decrease or loss of function, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide to non-target sequences under conditions in which specific binding is desired, *i.e.*, under physiological conditions in the case of *in vivo* assays or therapeutic treatment, or in the case of *in vitro* assays, under conditions in which the assays are performed.

Antisense oligonucleotides are commonly used as research reagents, diagnostic aids, and therapeutic agents. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes, for example to distinguish between the functions of various members of a biological pathway. This specific inhibitory effect has, therefore, been harnessed by those skilled in the art for research uses. Antisense oligonucleotides have also been used as diagnostic aids based on their specific binding or hybridization to DNA or mRNA that are present in certain disease states and due to the high degree of sensitivity that hybridization based assays and amplified assays that utilize some of polymerase chain reaction afford. The specificity and sensitivity of oligonucleotides is also harnessed by those of skill in the art for therapeutic uses. For example, the following U.S. patents demonstrate palliative, therapeutic and other methods utilizing antisense oligonucleotides. U. S. Patent No. 5,135,917 provides antisense oligonucleotides that inhibit human interleukin-1 receptor expression. U.S. Patent No. 5,098,890 is directed to antisense oligonucleotides complementary to the *c-myc* oncogene and antisense oligonucleotide therapies for certain cancerous

conditions. U.S. Patent No. 5,087,617 provides methods for treating cancer patients with antisense oligonucleotides. U.S. Patent No. 5,166,195 provides oligonucleotide inhibitors of Human Immunodeficiency Virus (HIV). U.S. Patent No. 5,004,810 provides oligomers capable of hybridizing to herpes simplex virus Vmw65 mRNA and inhibiting replication. U.S. Patent No. 5,194,428 provides antisense oligonucleotides having antiviral activity against influenza virus. U.S. Patent No. 4,806,463 provides antisense oligonucleotides and methods using them to inhibit HTLV-III replication. U.S. Patent No. 5,286,717 provides oligonucleotides having a complementary base sequence to a portion of an oncogene. U.S. Patent No. 5,276,019 and U.S. Patent No. 5,264,423 are directed to phosphorothioate oligonucleotide analogs used to prevent replication of foreign nucleic acids in cells. U.S. Patent No. 4,689,320 is directed to antisense oligonucleotides as antiviral agents specific to cytomegalovirus (CMV). U.S. Patent No. 5,098,890 provides oligonucleotides complementary to at least a portion of the mRNA transcript of the human *c-myc* gene. U.S. Patent No. 5,242,906 provides antisense oligonucleotides useful in the treatment of latent Epstein-Barr virus (EBV) infections. Other examples of antisense oligonucleotides are provided herein.

Specific examples of some preferred modified oligonucleotides envisioned for use in the ligand conjugated oligonucleotides of the present invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones or internucleoside linkages include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their intersugar backbone can also be considered to be oligonucleosides.

Specific oligonucleotide chemical modifications are described below. It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the following modifications may be incorporated in a single antisense compound or even in a single residue thereof, for example, at a single nucleoside within an oligonucleotide.

Preferred modified internucleoside linkages or backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

Representative United States Patents that teach the preparation of the above phosphorus atom containing linkages include, but are not limited to, U.S. Patents Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; and 5,697,248, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified internucleoside linkages or backbones that do not include a phosphorus atom therein (i.e., oligonucleosides) have backbones that are formed by short chain alkyl or cycloalkyl intersugar linkages, mixed heteroatom and alkyl or cycloalkyl intersugar linkages, or one or more short

chain heteroatomic or heterocyclic intersugar linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and
5 thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component
10 parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Patents Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564;
15 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of which are commonly owned with this application, and each of
20 which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleoside units are replaced with novel groups. The nucleobase units are maintained for hybridization with an
25 appropriate nucleic acid target compound. One such oligonucleotide, an oligonucleotide mimetic, that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide
30 containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to,

U.S. Patents Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497.

- 5 Some preferred embodiments of the present invention may employ oligonucleotides with phosphorothioate linkages and oligonucleosides with heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MMI backbone],
- 10 -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as -O-P-O-CH₂-] of the above referenced U.S. Patent 5,489,677, and the amide backbones of the above referenced U.S. Patent No. 5,602,240. Also preferred are oligonucleotides having
- 15 morpholino backbone structures of the above-referenced U.S. Patent No. 5,034,506.

The oligonucleotides employed in the ligand conjugated oligonucleotides of the present invention may additionally or alternatively comprise nucleobase (often referred to in the art

20 simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (a) and guanine (G), and the pyrimidine bases thymine (T), cytosine © and uracil (U). Modified nucleobases include other synthetic and natural nucleobases

25 such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-

30 propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines,

7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those
5 disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and*
10 *Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligonucleotides of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted
15 purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2 °C (*Id.*, pages 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-
20 methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Patent 3,687,808, as well as
25 U.S. Patents 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned, and each of which is herein incorporated by
30 reference, and commonly owned United States patent application 08/762,488, filed on December 10, 1996, also herein incorporated by reference.

The oligonucleotides employed in the ligand conjugated oligonucleotides of the present invention may additionally or

alternatively comprise one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl, O-, S-, or N-alkenyl, or O, S- or N-alkynyl, wherein the alkyl, alkenyl and alkynyl
5 may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following
10 at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, poly-alkylamino, substituted silyl, an RNA cleaving group, a
15 reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. a preferred modification includes 2'-methoxyethoxy
20 [2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE] (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486), i.e., an alkoxyalkoxy group. a further preferred modification includes 2'-dimethylaminooxyethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in co-owned United States
25 patent application Serial Number 09/016,520, filed on January 30, 1998, the contents of which are herein incorporated by reference.


Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F).
30 Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide.

As used herein, the term "sugar substituent group" or "2'-


substituent group" includes groups attached to the 2'-position of the ribofuranosyl moiety with or without an oxygen atom. Sugar substituent groups amenable to the present invention include, but are not limited to, fluoro, O-alkyl, O-alkylamino, O-alkylalkoxy, protected O-alkylamino, O-alkylaminoalkyl, O-alkyl imidazole and polyethers of the formula (O-alkyl)_m, wherein m is 1 to about 10. Preferred among these polyethers are linear and cyclic polyethylene glycols (PEGs), and (PEG)-containing groups, such as crown ethers and those which are disclosed by Ouchi et al. (*Drug Design and Discovery* **1992**, 9:93); Ravasio et al. (*J. Org. Chem.* **1991**, 56:4329); and Delgado et. al. (*Critical Reviews in Therapeutic Drug Carrier Systems* **1992**, 9:249), each of which is hereby incorporated by reference in its entirety. Further sugar modifications are disclosed by Cook (*Anti-Cancer Drug Design*, **1991**, 6:585-607). Fluoro, O-alkyl, O-alkylamino, O-alkyl imidazole, O-alkylaminoalkyl, and alkyl amino substitution is described in United States Patent Application serial number 08/398,901, filed March 6, 1995, entitled "Oligomeric Compounds having Pyrimidine Nucleotide(s) with 2' and 5' Substitutions," hereby incorporated by reference in its entirety.

Additional sugar substituent groups amenable to the present invention include 2'-SR and 2'-NR₂ groups, wherein each R is, independently, hydrogen, a protecting group or substituted or unsubstituted alkyl, alkenyl, or alkynyl. 2'-SR nucleosides are disclosed in United States Patent No. 5,670,633, issued September 23, 1997, hereby incorporated by reference in its entirety. The incorporation of 2'-SR monomer synthons is disclosed by Hamm et al. (*J. Org. Chem.*, **1997**, 62:3415-3420). 2'-NR nucleosides are disclosed by Goettingen, M., *J. Org. Chem.*, **1996**, 61, 6273-6281; and Polushin et al., *Tetrahedron Lett.*, **1996**, 37, 3227-3230. Further representative 2'-substituent groups amenable to the present invention include

those having one of formula XI or XII:



XI



XII

wherein:

E is C₁-C₁₀ alkyl, N(Q₃)(Q₄) or N=C(Q₃)(Q₄);

5 each Q₃ and Q₄ is, independently, H, C₁-C₁₀ alkyl, dialkylaminoalkyl, a nitrogen protecting group, a tethered or untethered conjugate group, a linker to a solid support;

or Q₃ and Q₄, together, form a nitrogen protecting group or a ring structure optionally including at least one
10 additional heteroatom selected from N and O;

q¹ is an integer from 1 to 10;

q² is an integer from 1 to 10;

q³ is 0 or 1;

q⁴ is 0, 1 or 2;

15 each Z₁, Z₂ and Z₃ is, independently, C₄-C₇ cycloalkyl, C₅-C₁₄ aryl or C₃-C₁₅ heterocyclyl, wherein the heteroatom in said heterocyclyl group is selected from oxygen, nitrogen and sulfur;

Z₄ is OM₁, SM₁, or N(M₁)₂;

20 each M₁ is, independently, H, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C(=NH)N(H)M₂, C(=O)N(H)M₂ or OC(=O)N(H)M₂;

M₂ is H or C₁-C₈ alkyl; and

Z₅ is C₁-C₁₀ alkyl, C₁-C₁₀ haloalkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₆-C₁₄ aryl, N(Q₃)(Q₄), OQ₃, halo, SQ₃ or CN.

25 Representative 2'-O-sugar substituent groups of formula XI are disclosed in United States Patent Application serial

number 09/130,973, filed August 7, 1998, entitled "Capped 2'-Oxyethoxy Oligonucleotides," hereby incorporated by reference in its entirety.

Representative cyclic 2'-O-sugar substituent groups of
5 formula XII are disclosed in United States Patent Application serial number 09/123,108, filed July 27, 1998, entitled "RNA Targeted 2'-Modified Oligonucleotides that are Conformationally Preorganized," hereby incorporated by reference in its entirety.

10 Sugars having O-substitutions on the ribosyl ring are also amenable to the present invention. Representative substitutions for ring O include, but are not limited to, S, CH₂, CHF, and CF₂. See, e.g., Secrist et al., *Abstract 21, Program & Abstracts, Tenth International Roundtable,*
15 *Nucleosides, Nucleotides and their Biological Applications,* Park City, Utah, Sept. 16-20, 1992, hereby incorporated by reference in its entirety.

Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar.
20 Representative United States patents that teach the preparation of such modified sugars structures include, but are not limited to, U.S. Patents Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909;
25 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned, and each of which is herein incorporated by reference, and commonly owned United States patent application 08/468,037, filed on June 5, 1995, also herein incorporated by reference.

30 Additional modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide and the 5' position of 5' terminal nucleotide. For example, one additional modification of the ligand conjugated oligonucleotides of the

present invention involves chemically linking to the oligonucleotide one or more additional non-ligand moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties
5 include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660,
10 306; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 111; Kabanov et al., *FEBS Lett.*, 1990, 259, 327; Svinarchuk et al.,
15 *Biochimie*, 1993, 75, 49), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan et al.,
20 *Nucleosides & Nucleotides*, 1995, 14, 969), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*,
25 1996, 277, 923).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Patents Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538;
30 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335;

4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136;
5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469;
5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098;
5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475;
5 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142;
5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923;
5,599,928 and 5,688,941, certain of which are commonly owned,
and each of which is herein incorporated by reference.

The present invention also includes compositions employing
10 antisense compounds which are chimeric compounds. "Chimeric"
antisense compounds or "chimeras," in the context of this
invention, are antisense compounds, particularly
oligonucleotides, which contain two or more chemically distinct
regions, each made up of at least one monomer unit, i.e., a
15 nucleotide in the case of an oligonucleotide compound. These
oligonucleotides typically contain at least one region wherein
the oligonucleotide is modified so as to confer upon the
oligonucleotide increased resistance to nuclease degradation,
increased cellular uptake, and/or increased binding affinity
20 for the target nucleic acid. An additional region of the
oligonucleotide may serve as a substrate for enzymes capable
of cleaving RNA:DNA or RNA:RNA hybrids. By way of example,
RNase H is a cellular endonuclease which cleaves the RNA strand
of an RNA:DNA duplex. Activation of RNase H, therefore,
25 results in cleavage of the RNA target, thereby greatly
enhancing the efficiency of oligonucleotide inhibition of gene
expression. Consequently, comparable results can often be
obtained with shorter oligonucleotides when chimeric
oligonucleotides are used, compared to phosphorothioate
30 oligodeoxynucleotides hybridizing to the same target region.
Cleavage of the RNA target can be routinely detected by gel
electrophoresis and, if necessary, associated nucleic acid
hybridization techniques known in the art. RNase H-mediated
target cleavage is distinct from the use of ribozymes to cleave

nucleic acids, and ribozymes are not comprehended by the present invention.

By way of example, such "chimeras" may be "gapmers," i.e., oligonucleotides in which a central portion (the "gap") of the oligonucleotide serves as a substrate for, e.g., RNase H, and the 5' and 3' portions (the "wings") are modified in such a fashion so as to have greater affinity for, or stability when duplexed with, the target RNA molecule but are unable to support nuclease activity (e.g., 2'-fluoro- or 2'-methoxyethoxy-substituted). Other chimeras include "hemimers," that is, oligonucleotides in which the 5' portion of the oligonucleotide serves as a substrate for, e.g., RNase H, whereas the 3' portion is modified in such a fashion so as to have greater affinity for, or stability when duplexed with, the target RNA molecule but is unable to support nuclease activity (e.g., 2'-fluoro- or 2'-methoxyethoxy-substituted), or vice-versa.

A number of chemical modifications to oligonucleotides that confer greater oligonucleotide:RNA duplex stability have been described by Freier et al. (*Nucl. Acids Res.*, 1997, 25, 4429). Such modifications are preferred for the RNase H-refractory portions of chimeric oligonucleotides and may generally be used to enhance the affinity of an antisense compound for a target RNA.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Patents Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are

commonly owned, and each of which is herein incorporated by reference, and commonly owned and allowed United States patent application serial number 08/465,880, filed on June 6, 1995, also herein incorporated by reference.

5 The present invention also includes compositions employing oligonucleotides that are substantially chirally pure with regard to particular positions within the oligonucleotides. Examples of substantially chirally pure oligonucleotides include, but are not limited to, those having
10 phosphorothioate linkages that are at least 75% Sp or Rp (Cook et al., U.S. Patent No. 5,587,361) and those having substantially chirally pure (Sp or Rp) alkylphosphonate, phosphoramidate or phosphotriester linkages (Cook, U.S. Patents Nos. 5,212,295 and 5,521,302).

15 The present invention further encompasses ligand conjugated oligonucleotides employing ribozymes. Synthetic RNA molecules and derivatives thereof that catalyze highly specific endoribonuclease activities are known as ribozymes. (See, generally, U.S. Patent No. 5,543,508 to Haseloff et al., issued
20 August 6, 1996, and U.S. Patent No. 5,545,729 to Goodchild et al., issued August 13, 1996.) The cleavage reactions are catalyzed by the RNA molecules themselves. In naturally occurring RNA molecules, the sites of self-catalyzed cleavage are located within highly conserved regions of RNA secondary
25 structure (Buzayan et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1986, 83, 8859; Forster et al., *Cell*, 1987, 50, 9). Naturally occurring autocatalytic RNA molecules have been modified to generate ribozymes which can be targeted to a particular cellular or pathogenic RNA molecule with a high degree of
30 specificity. Thus, ribozymes serve the same general purpose as antisense oligonucleotides (i.e., modulation of expression of a specific gene) and, like oligonucleotides, are nucleic acids possessing significant portions of single-strandedness. That is, ribozymes have substantial chemical and functional

identity with oligonucleotides and are thus considered to be equivalents for purposes of the present invention.

The oligonucleotides used in the conjugates of the present invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is also known to use similar techniques to prepare other oligonucleotides such as the phosphorothioates and alkylated derivatives.

Teachings regarding the synthesis of particular modified oligonucleotides may be found in the following U.S. patents or pending patent applications, each of which is commonly assigned with this application: U.S. Patents Nos. 5,138,045 and 5,218,105, drawn to polyamine conjugated oligonucleotides; U.S. Patent No. 5,212,295, drawn to monomers for the preparation of oligonucleotides having chiral phosphorus linkages; U.S. Patents Nos. 5,378,825 and 5,541,307, drawn to oligonucleotides having modified backbones; U.S. Patent No. 5,386,023, drawn to backbone modified oligonucleotides and the preparation thereof through reductive coupling; U.S. Patent No. 5,457,191, drawn to modified nucleobases based on the 3-deazapurine ring system and methods of synthesis thereof; U.S. Patent No. 5,459,255, drawn to modified nucleobases based on N-2 substituted purines; U.S. Patent No. 5,521,302, drawn to processes for preparing oligonucleotides having chiral phosphorus linkages; U.S. Patent No. 5,539,082, drawn to peptide nucleic acids; U.S. Patent No. 5,554,746, drawn to oligonucleotides having β -lactam backbones; U.S. Patent No. 5,571,902, drawn to methods and materials for the synthesis of oligonucleotides; U.S. Patent No. 5,578,718, drawn to nucleosides having alkylthio groups, wherein such groups may be used as linkers to other moieties attached at any of a variety of positions of the nucleoside; U.S. Patents Nos.

5,587,361 and 5,599,797, drawn to oligonucleotides having phosphorothioate linkages of high chiral purity; U.S. Patent No. 5,506,351, drawn to processes for the preparation of 2'-O-alkyl guanosine and related compounds, including 2,6-
5 diaminopurine compounds; U.S. Patent No. 5,587,469, drawn to oligonucleotides having N-2 substituted purines; U.S. Patent No. 5,587,470, drawn to oligonucleotides having 3-deazapurines; U.S. Patents Nos. 5,223,168, issued June 29, 1993, and 5,608,046, both drawn to conjugated 4'-desmethyl nucleoside
10 analogs; U.S. Patent Nos. 5,602,240, and 5,610,289, drawn to backbone modified oligonucleotide analogs; and U.S. patent application Serial No. 08/383,666, filed February 3, 1995, and U.S. Patent No. 5,459,255, drawn to, *inter alia*, methods of synthesizing 2'-fluoro-oligonucleotides.

15 In the ligand conjugated oligonucleotides and ligand molecule-bearing sequence-specific linked nucleosides of the present invention, the oligonucleotides and oligonucleosides may be assembled on a suitable DNA synthesizer utilizing standard nucleotide or nucleoside precursors, or nucleotide or
20 nucleoside conjugate precursors that already bear the linking moiety, or ligand-nucleotide or nucleoside conjugate precursors that already bear the ligand molecule.

When using nucleotide conjugate precursors that already bear a linking moiety, the synthesis of the sequence-specific
25 linked nucleosides is typically completed, and the ligand molecule is then reacted with the linking moiety to form the ligand conjugated oligonucleotide. This approach to the synthesis of oligonucleotide conjugates bearing a variety of molecules such as steroids, vitamins, lipids and reporter
30 molecules has previously been described (see Manoharan et al., PCT Application WO 93/07883). In a preferred embodiment the oligonucleotides or linked nucleosides of the present invention are synthesized by an automated synthesizer using phosphoramidites derived from ligand-nucleoside conjugates in addition

to the standard phosphoramidites and non-standard phosphoramidites that are commercially available and routinely used in oligonucleotide synthesis.

In application serial No. US91/00243, application serial
5 No. 463,358 and application serial No. 566,977, all incorporated herein by reference, it is reported that incorporation of a 2'-O-methyl, 2'-O-ethyl, 2'-O-propyl, 2'-O-allyl, 2'-O-aminoalkyl or 2'-deoxy-2'-fluoro group in nucleosides of an oligonucleotide confers enhanced hybridization
10 properties to the oligonucleotide. It is further reported that oligonucleotides containing phosphorothioate backbones have enhanced nuclease stability. Thus, functionalized, linked nucleosides of the invention can be augmented to further include either or both a phosphorothioate backbone or a 2'-O-
15 methyl, 2'-O-ethyl, 2'-O-propyl, 2'-O-aminoalkyl, 2'-O-allyl or 2'-deoxy-2'-fluoro group thereon.

In some preferred embodiments, functionalized nucleoside sequences of the invention possessing an amino group at the 5'-terminus are prepared using a DNA synthesizer, and then reacted
20 with an active ester derivative of a selected ligand. Active ester derivatives are well known to those skilled in the art. Representative active esters include N-hydrosuccinimide esters, tetrafluorophenolic esters, pentafluorophenolic esters and pentachlorophenolic esters. The reaction of the amino group
25 and the active ester produces an oligonucleotide in which the selected ligand is attached to the 5'-position through a linking group. The amino group at the 5'-terminus can conveniently be prepared utilizing the above noted 5'-Amino-Modifier C6 reagent. In a preferred embodiment, ligand
30 molecules may be conjugated to oligonucleotides at the 5'-position by the use of a ligand-nucleoside phosphoramidite wherein the ligand is linked to the 5'-hydroxy group directly or indirectly via a linker. Such ligand-nucleoside phosphoramidites are typically used at the end of an automated

synthesis procedure to provide a ligand conjugated oligonucleotide bearing the ligand at the 5'-terminus.

In further preferred embodiments, functionalized nucleoside sequences of the invention can be prepared wherein
5 a selected ligand is attached to the 3'-terminal amino group using a 3'-amino modified controlled pore glass (sold by Clontech Laboratories Inc., Palo Alto, CA) and subsequent attachment of the ligand is achieved by reaction with a ligand active ester.

10 In another preferred embodiment of the present invention, the ligand may be attached to the oligonucleotide at the 3'-terminus through the use of an appropriate multifunctional linker such as a universal linker. In this case the ligand is first derivatized with the universal linker
15 and this conjugate then loaded onto a solid support. Subsequent synthesis of nucleic acids or oligonucleotides on this solid support affords upon cleavage and deprotection the ligand conjugated oligonucleotide bearing a ligand molecule at the 3'-terminus.

20 In still further preferred embodiments, functionalized sequences of nucleosides and ligand conjugated oligonucleotides of the present invention can be prepared wherein the ligand molecule is attached either directly or via a linking group to any one of the atoms of the nucleobase of any of the nucleoside
25 units of the oligonucleotide. Thus, one or more ligand molecules may be attached to the nucleobase at the 3'-terminus, the 5'-terminus or any position in between. Such attachment can be accomplished, for example, by chemistries described in the literature, and mentioned above. The preferred mode of
30 attachment of ligand molecules to nucleobases is via the intermediacy of an appropriate linker present on a nucleoside precursor. The ligand-nucleoside conjugate is then phosphitylated at the 3'-position to afford a ligand-nucleoside conjugate phosphoramidite which may be used subsequently as a

building block together with traditional nucleoside phosphoramidites for the automated synthesis of oligonucleotides. The number and location of insertion of such ligand nucleotide conjugate phosphoramidites will then dictate
5 the number and location of ligand molecules present in the synthesized ligand conjugated oligonucleotide of the present invention.

The present invention also provides ligand conjugated oligonucleotides wherein the ligand molecule is attached to one
10 of the atoms of the internucleotide linkage. One typical internucleotide linkage in nucleic acids and oligonucleotides is the phosphodiester linkage. Numerous modified internucleotide linkages are known in the art including, but not limited to, phosphorothioate, methyl phosphonate, and
15 phosphordithioate, as described above. Ligand molecules may be conjugated at one of the atoms of such internucleotide linkages with or without the intermediacy of a linking group. Attachment of the ligand molecule may be accomplished in accordance with the methods of the invention either during the
20 preparation of the nucleoside building block such as the phosphoramidite or may be performed during the formation of the internucleotide linkage during oligonucleotide synthesis.

In further preferred embodiments of the invention, the ligand molecule is attached at multiple sites on one
25 oligonucleotide. For example, ligand conjugated oligonucleotides can be prepared wherein one or more ligands are attached to both ends of a linked nucleoside sequence. Preferably such a structure is prepared by reacting a 3',5'-diamino sequence with a ligand active ester. The required
30 oligonucleoside sequence can be synthesized, for example, utilizing the 3'-Amino-Modifier and the 5'-Amino-Modifier C6 (or Aminolink-2) reagents noted above or by utilizing the above noted 3'-amino modified controlled pore glass reagent in combination with the 5'-Amino-Modifier C2 (or Aminolink-2)

reagents. Alternatively, such multiply conjugated oligonucleotides may readily be synthesized according to the methods of the invention using an appropriate ligand-nucleoside conjugate phosphoramidites as and where needed in a given 5 oligonucleotide sequence during automated synthesis.

In still further preferred embodiments of the invention, an oligonucleoside sequence bearing an aminolinker at the 2'-position of one or more selected nucleosides is prepared using a suitably functionalized nucleotide such as, for example, 5'-10 dimethoxytrityl-2'-O-(ϵ -phthalimidylaminopentyl)-2'-deoxyadenosine-3'-N,N-diisopropyl-cyanoethoxy phosphoramidite. See the above referenced patent applications serial numbers US91/00243, 566,977 and 463,358. Preferably, the nucleotide or nucleotides are attached to the ligand by reaction with an 15 active ester or a thioisocyanate thereof, at one or more of the nucleoside components of the oligonucleotide.

In yet further preferred embodiments, functionalized nucleoside sequences of the invention can be prepared wherein the heterocyclic base of one or more nucleosides can be linked 20 to a ligand molecule. for example, utilizing 5'-O-dimethoxytrityl-5-[N(7-trifluoroacetyl aminoheptyl)-3-acrylamido]-2'-deoxyuridine 3'-O-(methyl N,N-diisopropyl)phosphoramidate as described by Jablonski et. al. *supra* (also commercially available from Glen Research) the desired 25 nucleoside, functionalized to incorporate a linking group on its heterocyclic base, is incorporated into the linked nucleoside sequence using a DNA synthesizer.

In further functionalized linked nucleosides of the invention, conjugation (or linking) of ligand molecules is 30 achieved by conjugation of the ligand to the above described amino linking group on the nucleoside. This can be effected in several ways. For example, a ligand-nucleoside conjugate of the invention can be prepared by conjugation of the ligand molecule to the nucleoside using EDC/sulfo-NHS (i.e. 1-ethyl-

3(3-dimethylaminopropylcarbodiimide/N-hydroxysulfosuccinimide) to conjugate the carboxylate function of the ligand with the amino function of the linking group on the nucleoside.

Ligand conjugated oligonucleotides of the present
5 invention may be prepared by conjugation of the ligand molecule to the nucleoside sequence via a heterobifunctional linker such as *m*-maleimidobenzoyl-N-hydroxysulfosuccinimide ester (MBS) or succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC), to link a nucleophilic position, preferably a thiol,
10 on the ligand molecule to the amino function of the linking group on nucleoside sequence. By this mechanism, an oligonucleoside-maleimide conjugate is formed by reaction of the amino group of the linker on the linked nucleosides with the MBS or SMCC maleimide linker. The conjugate is then
15 reacted with ligand molecules, preferably those that possess a thiol functionality.

Alternatively, an ligand conjugated oligonucleotide can be prepared by conjugation of the ligand molecule to the oligonucleotide or nucleoside via a homobifunctional linker
20 such as disuccinimidyl suberate (DSS), to link an amino function on the ligand to the amino group of a linker on the oligonucleotide sequence. By this mechanism, an oligonucleoside-succinimidyl conjugate is formed by reaction of the amino group of the linker on the nucleoside sequence
25 with a disuccinimidyl suberate linker. The disuccinimidyl suberate linker couples with the amine linker on the nucleoside to extend the size of the linker. The extended linker is then reacted with an amino group of the ligand molecule.

A number of non-ligand molecules have been conjugated to
30 oligonucleotides in order to enhance the activity, cellular distribution or cellular uptake of the oligonucleotide, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties such as cholesterol (Letsinger et al.,

Proc. Natl. Acad. Sci. USA, 1989, 86:6553), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4:1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660:306; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3:2765), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20:533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10:111; Kabanov et al., *FEBS Lett.*, 1990, 259:327; Svinarchuk et al., *Biochimie*, 1993, 75:49), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651; Shea et al., *Nucl. Acids Res.*, 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14:969), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277:923).

Representative United States patents that teach the preparation of such oligonucleotide conjugates have been listed above. Typical conjugation protocols involve the synthesis of oligonucleotides bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction may be performed either with the oligonucleotide still bound to the solid support or following cleavage of the oligonucleotide in solution phase. Purification of the oligonucleotide conjugate by HPLC typically affords the pure conjugate.

Alternatively, the molecule being conjugated may be converted into a building block such as a phosphoramidite via an alcohol group present in the molecule or by attachment of


a linker bearing an alcohol group that may be phosphitylated.

Each of these approaches may be used for the synthesis of ligand conjugated oligonucleotides. Aminolinked oligonucleotides may be coupled directly with ligand via the use of coupling reagents or following activation of the ligand as an NHS or pentfluorophenolate ester. Ligand phosphoramidites may be synthesized via the attachment of an aminohexanol linker to one of the carboxyl groups followed by phosphitylation of the terminal alcohol functionality. Other linkers such as cysteamine may also be utilized for conjugation to a chloroacetyl linker present on a synthesized oligonucleotide.

In one preferred embodiment of the methods of the invention, the preparation of ligand conjugated oligonucleotides commences with the selection of appropriate precursor molecules upon which to construct the ligand molecule. Typically the precursor is an appropriately protected derivative of the commonly used nucleosides. For example, the synthetic precursors for the synthesis of the ligand conjugated oligonucleotides of the present invention include, but are not limited to, 2'-6-aminoalkoxy-5'-ODMT-nucleosides, 2'-6-aminoalkylamino-5'-ODMT-nucleosides, 5'-6-aminoalkoxy-2'-deoxynucleosides, 5'-6-aminoalkoxy-2'-protected-nucleosides, 3'-6-aminoalkoxy-5'-ODMT-nucleosides, and 3'-aminoalkylamino-5'-ODMT-nucleosides that may be further protected in the nucleobase portion of the molecule. The use of such precursors is anticipated to afford ligand conjugated oligonucleotides where attachment is at one of many possible sites such as the 2', 3' or 5' position of one or more of the nucleoside components of oligonucleotides. Methods for the synthesis of such aminolinked protected nucleoside precursors are known to the art skilled and are available in the literature.

In one embodiment of the invention a conjugated oligonucleotide is prepared starting with a conjugated

nucleoside using solid phase chemistries. An oligonucleoside is selected having a serum protein binding ligand attached or optionally linked to one the 2', 3', or 5' positions, a protected hydroxyl at one of the 2', 3', or 5' positions and a free hydroxyl group located at the other one of the 2', 3', or 5' positions. The free hydroxyl group is treated with a bi-functional linking moiety and the resulting nucleoside is reacted with a solid support. A representative conjugated nucleoside attached via a succinyl linker at the 2'-O-position to a solid support (from Example 20) is shown below:



The ligand is ibuprofen and the linker is a preferred 6-aminoheptyloxy linking group. Bx is a heterocyclic base moiety and Pg is a hydroxyl protecting group. The resulting solid support bound conjugated nucleoside is treated with a weak acid to remove the hydroxyl protecting group and treated with a further nucleoside or nucleotide to form a dimer. In one aspect of the invention the coupling of further nucleosides to form a desired oligonucleotide is performed using phosphoramidite monomers following known methods and procedures.

As used herein, the term "alkyl" includes but is not limited to straight chain, branch chain, and alicyclic hydrocarbon groups. Alkyl groups of the present invention may be substituted. Representative alkyl substituents are disclosed in United States Patent No. 5,212,295, at column 12,

lines 41-50, hereby incorporated by reference in its entirety.

As used herein, the term "aralkyl" denotes alkyl groups which bear aryl groups, for example, benzyl groups. The term "alkaryl" denotes aryl groups which bear alkyl groups, for example, methylphenyl groups. "Aryl" groups are aromatic cyclic compounds including but not limited to substituted and unsubstituted aromatic hydrocarbyl groups. Aralkyl groups (generally C₇-C₂₀) include but are not limited to groups having both aryl and alkyl functionalities, such as benzyl and xylyl groups. Preferred aryl and aralkyl groups include, but are not limited to, phenyl, benzyl, xylyl, naphthyl, toluyl, pyrenyl, anthracyl, phenanthryl, azulyl, phenethyl, cinnamyl, benzhydryl, and mesityl. Typical substituents for substitution include, but are not limited to, hydroxyl, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, or alkyl, aryl, alkenyl, or alkynyl groups.

As used herein, the term "alkanoyl" has its accustomed meaning as a group of formula -C(=O)-alkyl. A preferred alkanoyl group is the acetoxy group.

In general, the term "hetero" denotes an atom other than carbon, preferably but not exclusively N, O, or S, SO and SO₂. Accordingly, the term "heterocycle" denotes a cyclic structure having at least one non-carbon atom. "Cyclo" or "cyclyl" includes a cyclic group which may be mono-, bi- or tricyclic, and may be substituted with substituents such as oxo, acyl, alkoxy, alkoxycarbonyl, alkyl, alkenyl, alkynyl, amino, amido, azido, aryl, heteroaryl, carboxylic acid, cyano, guanidino, halo, haloalkyl, haloalkoxy, hydrazino, ODMT, alkylsulfonyl, nitro, sulfide, sulfone, sulfonamide, thiol and thioalkoxy.

Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples, which are not intended to be limiting.

Example 1**3'-O-Hexylaminofenbufenyl-5'-O-DMT-5-methyluridine (1)**

To a solution of 3'-O-(6-aminohexyl)-5-methyluridine (1.0 g, 1.51 mmol) (prepared according to the method described in Manoharan et al. (*Tetrahedron Lett.*, **1995**, 36:3647) dissolved in CH₂Cl₂ (15 mL) was added fenbufen (sigma, 424 mg, 1.66 mmol), followed by *N,N'*-dicyclohexylcarbodiimide (Fluka, 342 mg, 1.66 mmol) with shaking for about 2 hours. The mixture was filtered to remove dicyclohexylurea and the filtrate was partitioned between CH₂Cl₂ (50 mL) and saturated sodium bicarbonate solution (50 mL). The organic layer was dried over anhydrous sodium sulfate and evaporated. The resultant foam was purified by silica gel column chromatography using 50:50 EtOAc:hexanes as the eluent to give 1.75 g (92%) of the title compound as a colorless solid.

¹H NMR (CDCl₃): δ (m, 4H), 2.62-2.66 (m, 2H), 3.17-3.56 (m, 10H), 3.76-3.78 (m, 6H, OMe), 4.03-4.32 (m, 2H), 5.38-5.42 (d, 1H), 5.93-5.94 (d, 1H), 6.11 (t, 1H), 6.81-8.06 (m, aromatic), 9.6 (6, 1H, NH). ¹³C NMR (CDCl₃): δ 14.15, 20.99, 25.58, 26.41, 29.40, 30.27, 34.13, 39.35, 55.21, 62.38, 70.75, 73.84, 81.34, 87.98, 89.64, 102.39, 113.26, 126.13, 127.17-135.214 (m), 139.72, 140.123, 144.212, 145.83, 150.59, 158.64, 163.21, 172.15, 198.85.

Example 2**3'-O-Hexylaminofenbufenyl-2'-O-succinate-5'-O-DMT-5-methyluridine (2)**

Compound 1 (1.00 g, 1.12 mmol), succinic anhydride (0.168 g, 1.68 mmol), dimethylaminopyridine (0.068 g, 0.56 mmol), and triethylamine (0.16 mL, 1.12 mmol) were dissolved in 1,2-dichloroethane (3 mL) at room temperature. The reaction mixture, in a test-tube with a screw cap top, was placed in a heating block at 55 °C for 2 hours and then allowed to cool to

room temperature overnight. TLC using EtOAc:MeOH (85/15; v/v) showed complete conversion of the starting material. 1,2-Dichloroethane (30 mL) was added and the mixture was washed three times with portions of cold 10% citric acid (17 mL, aq) followed by three washes with portions of water (17 mL). The organic-phase was dried over sodium sulfate and evaporated to 1.14 g (100%) of the title compound as a foam.

¹H NMR (DMSO-d₆): δ 12.3 (s, 1H), 11.44 (s, 1H), 8.08-6.89 (m, 23H), 5.84 (d, 1H), 5.47 (m, 1H), 5.40 (d, 1H), 4.24 (m, 1H), 3.98 (m, 1H), 3.90 (m, 2H), 3.75 (s, 3H), 3.73 (s, 3H), 3.27 (m, 7H), 3.0 (m, 2H), 2.59 (m, 7H), 1.30 (m, 8H). (Kumar et al., *Nucleosides & Nucleotides*, **1993**, 12:565-584).

Example 3

3'-O-Hexylaminofenbufenyl-2'-O-succinate-5'-O-DMT-5-methyluridine LCAA-CPG (3)

Compound 2 (1.04 g, 1.05 mmol) and 4-methylmorpholine (0.23 mL, 2.10 mmol) were dissolved in DMF (19 mL) at room temperature. 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetra-methyl-uronium tetrafluoroborate (0.34 g, 1.05 mmol) and acid washed LCAA-CPG (4.56 g, 0.52 mmol) were added and the mixture was shaken overnight. The resulting resin was then washed three times with CH₂Cl₂ and three times with ether. The initial loading was found to be 41 μmol/g. The resin was then combined with Cap A (20 mL) and Cap B (20 mL) solutions from PerSeptive Biosystems GmbH, and shaken for another hour and washed with three portions of CH₂Cl₂ and ether. The capped resin 3 was placed under vacuum to dry overnight and the loading was determined to be 46 μmol/g.

Example 4

3'-O-(6-Aminohexyl-ketoprofenyl)-5'-O-DMT-5'-methyluridine (4)

To a solution of 3'-O-(6-aminohexyl)-5-methyluridine

(1.0 g, 1.51 mmol) in CH_2Cl_2 (15 mL) was added a solution of ketoprofen (sigma, 422 mg, 1.66 mmol) and DCC (Fluka, 342 mg, 1.66 mmol) with shaking for 2 hrs. The mixture was filtered and the filtrate was partitioned between CH_2Cl_2 (50 mL) and a solution of saturated sodium bicarbonate (50 mL). The organic layer was separated, dried over anhydrous sodium sulfate and evaporated. The residual foam was purified by silica gel column chromatography using 50:50 ethylacetate:hexanes as the eluant to give 1.82 g (88%) of the title compound.

^{13}C NMR (CDCl_3): δ 18.63, 25.55, 26.38, 29.32, 39.44, 46.90, 55.20, 62.41, 70.75, 73.82, 81.33, 86.99, 89.53, 102.39, 113.24, 127.12, 127.99, 128.29, 129.65, 129.02, 130.02, 131.48, 132.56, 135.07, 135.19, 137.01, 137.89, 140.11, 142.04, 144.108, 150.53, 158.64, 163.11, 173.51.

Example 5

3'-O-Hexylaminoketoprofenyl-2'-O-succinate-5'-O-DMT-5-methyl-uridine (5)

Compound 4 (1.00 g, 1.12 mmol), succinic anhydride (0.168 g, 1.68 mmol), dimethylaminopyridine (0.068 g, 0.56 mmol), and triethylamine (0.16 mL, 1.12 mmol) were dissolved in 1,2-dichloroethane (3 mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 2 hours and then cooled to room temperature. TLC using EtOAc:MeOH (85/15; v/v) showed the absence of starting material. The mixture was diluted with 1,2-dichloroethane (30 mL) and washed three times with cold 10% citric acid (aqueous, 17 mL) and three times with water (17 mL). The organic-phase was dried over sodium sulfate and evaporated to give 1.14 g (100%) of the title compound as a foam.

^1H NMR ($\text{DMSO}-d_6$): δ 12.3 (s, 1H), 11.45 (s, 1H), 8.02 (m, 1H), 7.82-6.88 (m, 23H), 5.84 (d, 1H), 5.48 (m, 1H), 5.40

(d, 1H), 4.24 (m, 1H), 3.99 (m, 1H), 3.92 (m, 2H), 3.74 (s, 6H), 3.32 (m, 5H), 3.00 (m, 2H), 2.51 (m, 5H), 1.36 (m, 7H), 1.14 (s, 4H).

Example 6

5 3'-O-Hexylaminoketoprofenyl-2'-O-succinate-5'-O-DMT-5-methyluridine LCAA-CPG (6)

Compound 5 (1.04 g, 1.05 mmol) and 4-methylmorpholine (0.23 mL, 2.10 mmol) were dissolved in DMF (19 mL) at room temperature. 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (0.34 g, 1.05 mmol) and acid washed LCAA-CPG (4.56 g, 0.52 mmol) were added and the mixture was shaken overnight. The resulting resin was washed with CH₂Cl₂ (x3) and ether (x3). The initial loading was found to be 32 μmol/g. The resin was combined with Cap A (20 mL) and 15 Cap B (20 mL) solutions from PerSeptive Biosystems GmbH, and shaken for another hour. The resin was washed again with CH₂Cl₂ (x3) and ether (x3). The capped resin (6) was dried under vacuum overnight. The loading was determined to be 44 μmol/g.

20 Example 7

3'-O-(6-Aminohexyl-suprofenyl)-5'-O-DMT-5-methyluridine (7)

To a solution of 3'-O-(6-aminohexyl)-5-methyluridine (1.0 g, 1.51 mmol) in CH₂Cl₂ (15 mL) was added a solution of suprofen (sigma, 432 mg, 1.66 mmol) followed by DCC (Fluka, 342 mg, 1.66 mmol). After shaking the reaction mixture for 2 25 hours, dicyclohexyl urea was filtered off. The resulting organic solution was partitioned between CH₂Cl₂ (50 mL) and a solution of saturated NaHCO₃. The organic layer was separated, dried over anhydrous sodium sulfate and evaporated. The 30 resulting product was purified by silica gel column chromatography using 50:50 ethylacetate:hexanes as the eluant

to give 1.75 g (88%) of the title compound as a colorless solid.

¹³C NMR (CDCl₃): δ 14.06, 18.61, 25.56, 26.41, 29.33, 29.44, 39.49, 47.08, 55.21, 60.36, 62.43, 70.75, 73.83, 81.352, 87.00, 89.54, 102.39, 113.24, 127.13, 127.65, 127.98, 129.68, 130.03, 134.30, 134.82, 135.06, 135.19, 136.99, 141.122, 143.41, 114.17, 146.16, 150.52, 158.60, 163.24, 173.25, 187.696.

Example 8

10 **3'-O-Hexylaminosuprofenyl-2'-O-succinate-5'-O-DMT-5-methyluridine (8)**

Compound 7 (1.00 g, 1.11 mmol), succinic anhydride (0.167 g, 1.66 mmol), dimethylaminopyridine (0.068 g, 0.56 mmol), and triethylamine (0.15mL, 1.11 mmol) were dissolved in 1,2-dichloroethane (3mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 2 hours and then cooled to room temperature overnight. TLC using EtOAc:MeOH (85:15, v/v) showed that all the starting material had been converted. The mixture was diluted with 1,2-dichloroethane (30 mL) washed three times with cold 10% acid (aqueous, 17 mL) and three times with water (17 mL). The organic-phase was dried over sodium sulfate and evaporated to give 1.13 g (100%) of the title compound as a foam.

25 ¹H NMR (DMSO-d₆): δ 12.3 (s, 1H), 11.44 (s, 1H), 8.12-6.88 (m, 21H), 5.84 (d, 1H), 5.48 (m, 1H), 5.40 (d, 1H), 4.24 (m, 1H), 3.99 (m, 1H), 3.91 (m, 3H), 3.74 (s, 6H), 3.32 (m, 5H), 3.02 (m, 2H), 2.54 (m, 5H), 1.37 (m, 7H), 1.16 (s, 4H).

Example 9

30 **3'-O-Hexylaminosuprofenyl-2'-O-succinate-5'-O-DMT-5-**

methyluridine LCAA-CPG (9)

Compound 8 (1.03 g, 1.03 mmol) and 4-methylmorpholine (0.23mL, 2.06 mmol) were dissolved in DMF (19mL) at room temperature. 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (0.33 g, 1.03 mmol) and acid washed LCAA-CPG (4.47 g, 0.52 mmol) were added and the mixture was shaken overnight. The resulting resin was then washed with CH₂Cl₂ (x3) and ether (x3). The initial loading was found to be 36 µmol/g. The resin was then combined with
10 Cap A (20mL) and Cap B (20mL) solutions from PerSeptive Biosystems GmbH, and shaken for one hour. The resin was washed with CH₂Cl₂ (x3) and ether (x3). The capped resin 9 was dried under vacuum overnight and the loading was determined to be 47 µmol/g.

15 Example 10**3'-O-(6-Aminohexyl-carprofenyl)-5'-O-DMT-5-methyluridine (10)**

To a solution of 3'-O-(6-aminohexyl)-5-methyluridine (1.0 g, 1.51 mmol) in CH₂Cl₂ (15 mL) was added carprofen (sigma, 453 mg, 1.66 mmol) followed by DCC (Fluka, 342 mg, 1.66 mmol).
20 After shaking the reaction mixture for 2 hours, dicyclohexyl urea was filtered off. The resulting organic solution was partitioned between CH₂Cl₂ (50 mL) and a solution of saturated NaHCO₃. The organic layer was dried over anhydrous sodium sulfate, evaporated. The resulting product was purified by
25 silica gel column chromatography using 50:50 ethylacetate:hexanes as the eluant to give 1.65 g (84%) of the title compound as a colorless solid.

Example 11**3'-O-Hexylaminocarprofenyl-2'-O-succinate-5'-O-DMT-5-methyluridine (11)**
30

Compound 10 (1.00 g, 1.09 mmol), succinic anhydride

(0.164 g, 1.64 mmol), dimethylaminopyridine (0.066 g, 0.54 mmol), and triethylamine (0.15mL, 1.09 mmol) were dissolved in 1,2-dichloroethane (3 mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 2 hours and then cooled to room temperature. TLC using EtOAc:MeOH (85/15/ v/v) showed that the starting material was converted. The mixture was diluted with 1,2-dichloroethane (30 mL) and washed three times with cold 10% citric acid (aqueous, 17 mL) and three times with water (17 mL). The organic phase was dried over sodium sulfate and evaporated to give 1.07 g (97%) of the title compound as a foam.

¹H NMR (DMSO-d₆): δ 12.3 (s, 1H), 11.45 (s, 1H), 11.36 (s, 1H), 8.16-6.88 (m, 20H), 5.84 (d, 1H), 5.48 (m, 1H), 5.41 (d, 1H), 4.23 (m, 1H), 3.99 (m, 1H), 3.92 (m, 1H), 3.74 (s, 6H), 3.32 (m, 5H), 3.02 (m, 2H), 2.54 (m, 5H), 1.29 (m, 12H).

Example 12

3'-O-Hexylaminocarprofenyl-2'-O-succinate-5'-O-DMT-5-methyl-uridine LCAA-CPG (12)

Compound 11 (0.970 g, 0.96 mmol) and 4-methylmorpholine (0.21mL, 1.92 mmol) were dissolved in DMF (19mL) at room temperature. 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate (0.31 g, 0.96 mmol) and acid washed LCAA-CPG (4.14 g, 0.48 mmol) were added and the mixture was shaken overnight. The resulting resin was washed with CH₂Cl₂ (x3) and ether (x3). The initial loading was found to be 39 μmol/g. The resin was then combined with Cap A (20mL) and Cap B (20mL) solutions from PerSeptive Biosystems GmbH, and shaken for one hour. The resin was again washed with CH₂Cl₂ (x3) and ether (x3). The capped resin 12 was dried under vacuum. The loading was determined to be 41 μmol/g.

Example 13**3'-O-(6-Aminohexyl-palmityl)-5'-O-DMT-2'-O-succinyluridine (13)**

To a solution of 3'-O-(6-aminoethyl)-5'-O-DMT-uridine (1.5 g, 2.33 mmol) in CH₂Cl₂ (20 mL) at room temperature was added diisopropylamine (0.81 mL, 4.66 mmol) followed by palmitic acid pentafluorophenyl ester (compound, *vide infra*, 1.04 g, 2.8 mmol) with stirring overnight. The mixture was concentrated and the residue purified by silica gel column chromatography using EtOAc: CH₃OH (90:10; v/v) as the eluent to give 1.39 g (72%) of the title compound without the 2'-O-succinyl group attached.

The above conjugate (1.12 mmol), succinic anhydride (0.17g, 1.7 mmol), dimethyl amino pyridine (0.068 g, 0.56 mmol) and triethyl amine (0.16 mL, 1.12 mmol) were dissolved in 1,2-dichloroethane (3 mL) at room temperature in a test-tube with a screw cap top. The reaction mixture was placed in a heating block at 55 °C for 2 hours and then cooled to room temperature. TLC using EtOAc:MeOH (85:15; v/v) showed the absence of starting material. The mixture was diluted with 1,2-dichloroethane (30 mL) and washed three times with cold 10% aqueous citric acid (25 mL) and three times with water (25 mL). The organic phase was dried over sodium sulfate and evaporated to give 1.2 g (quantitative yield) of the title compound as a foam.

Example 14**5'-O-DMT-3'-O-palmitylaminoethyl-2'-O-succinyluridine LCAA-CPG (14)**

Compound 13 (1.35 g, 1.35 mmol) and 4-dimethylaminopyridine (0.16 g, 1.35 mmol) were dissolved in CH₃CN (12.2 mL) at room temperature in a first flask. In a second flask, 2,2'-dithiobis-5-nitropyridine (0.42 g, 1.35 mmol) was dissolved in acetonitrile (8.53 mL) and dichloromethane (3.64

mL) and the resulting solution was added to the first flask. In a third flask, triphenylphosphine (0.35 g, 1.35 mmol) was dissolved in anhydrous CH_3CN (12.2 mL) and the resulting solution was added to the first flask. Acid washed LCAA-CPG
5 (10.9 g, having a loading of 115 mol/g) was added and the mixture was shaken for about 3 hours. The resulting resin was washed with CH_3CN (x3) followed by CH_2Cl_2 and ether to removed excess reagents. To the washed resin was added acetic anhydride (25 mL) in tetrahydrofuran (THF) and 1-
10 methylimidazole (25 mL) in THF (Cap A and Cap B reagents from Perseptive Biosystems GmbH) and the mixture was shaken for an 2 hours. The resin was again washed again with CH_2Cl_2 (x3) and ether (x3). The washed resin was dried overnight in a vacuum oven at room temperature under P_2O_5 . The yield of dried resin
15 was 10.8 g with the loading determined to be 44 mol/g.

A portion of the final resin (3.8 mg) was cleaved by treatment with trichloroacetic acid (25 mL, 3%) in CH_2Cl_2 . The loading was determined by measuring the absorption of released trityl cation at 503 nm on a spectrophotometer (Hewlett packard
20 8452A Diode Array spectrophotometer). The final derivatized resin yield was 10.8 g total.

Example 15

3'-O-Hexylaminopalmityl-5'-O-DMT-cytidine (15)

To 3'-O-hexylamino-5'-O-DMT-cytidine (1.50 g, 2.33 mmol)
25 (purchased from RI Chemical, CA) dissolved in CH_2Cl_2 (20 mL) at room temperature was added diisopropylamine (0.81 mL, 4.66 mmol) and palmitic acid pentafluorophenyl-ester (1.18 g, 2.80 mmol) with stirring overnight. The mixture was was evaporated and the resulting crude purified by silica gel column
30 chromatography using EtOAc:MeOH (90/10; v/v) as the eluant to give 1.20 g (59%) of the title compound.

^1H NMR (CDCl_3): δ 7.99 (d, 1H), 7.40-6.81 (m, 15H), 5.91 (d, 1H), 5.63 (m, 1H), 5.49 (d, 1H), 4.32 (m, 1H), 4.19 (m, 1H),

4.03 (m, 1H), 3.79 (s, 6H), 3.54 (m, 2H), 3.38 (m, 2H), 3.20 (m, 2H), 2.14 (t, 2H), 1.58-1.24 (m, 34H), 0.87 (t, 3H). ^{13}C (CDCl_3): δ 173.32, 165.30, 158.63, 156.01, 144.40, 141.54, 135.48, 135.37, 130.09, 128.11, 127.96, 127.06, 113.25, 94.35, 91.36, 86.76, 81.37, 74.27, 70.85, 62.02, 56.30, 55.26, 52.82, 39.33, 36.86, 31.92, 29.69, 29.52, 29.36, 26.60, 25.85, 25.70, 22.69, 18.14, 14.11. MS (ES^-) calculated for $\text{C}_{52}\text{H}_{74}\text{N}_4\text{O}_8$ 882.6; Observed 881.8.

Example 16

10 3'-O-Hexylaminopalmityl-5'-O-DMT-N4-Benzoylcytidine (16)

To compound 15 (1.20 g, 1.36 mmol) dissolved in N,N-dimethylformamide (30 mL) at room temperature was added benzoic anhydride (0.37 g, 1.63 mmol) with stirring overnight. Saturated aqueous sodium bicarbonate was added and the mixture was extracted with ethyl acetate (x3). The organic-phase was dried over magnesium sulfate and evaporated. The crude product was then purified by silica gel column chromatography using EtOAc:MeOH (95/5; v/v) as the eluant to give 0.90 g (67%) of the title compound.

^1H NMR (CDCl_3) δ 8.75 (s, 1H), 8.41 (d, 1H), 7.92-6.85 (m, 20H), 6.00 (d, 1H), 5.49 (m, 1H), 4.38 (m, 1H), 4.27 (m, 1H), 4.09 (m, 1H), 3.82 (s, 6H), 3.50 (m, 5H), 3.23 (m, 2H), 2.15 (t, 2H), 1.45 (m, 34H), 0.88 (t, 3H); ^{13}C (CDCl_3) δ 173.15, 162.25, 158.72, 144.83, 144.07, 135.48, 135.28, 133.19, 130.10, 129.99, 129.05, 128.15, 128.05, 127.54, 127.20, 113.35, 91.80, 87.04, 81.79, 77.23, 76.97, 74.55, 71.00, 61.82, 55.26, 39.30, 36.91, 31.92, 29.69, 29.65, 29.50, 29.36, 26.60, 25.83, 25.72, 22.69, 14.12. MS (ES^-) calculated for $\text{C}_{59}\text{H}_{78}\text{N}_4\text{O}_9$ - DMT group 683.2. Observed 681.1 (16 without DMT group).

30 Example 17

3'-O-Hexylaminopalmityl-2'-O-succinate-5'-O-DMT-N4-benzoyl-

cytidine (17)

Compound 16 (0.88 g, 0.89 mmol), succinic anhydride (0.134 g, 1.34 mmol), dimethylaminopyridine (0.054 g, 0.44 mmol), and triethylamine (0.12 mL, 0.89 mmol) were dissolved in 1,2-dichloroethane (4mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 1 hour and cooled to room temperature overnight. TLC using EtOAc:MeOH (90/10; v/v) showed conversion of the starting material. The mixture was diluted with CH₂Cl₂ (40 mL) and the mixture was washed with cold 10% citric acid (20 mL, aq, x3) followed by water (20 mL, x3). The organic phase was dried over magnesium sulfate and evaporated to give 0.97 g (100%) of the title compound as a foam.

¹H NMR (DMSO-d₆) δ 8.34 (d, 1H), 7.95-6.86 (m, 20H), 6.10 (d, 1H), 5.72 (m, 1H), 5.37 (m, 1H), 4.18 (m, 2H), 3.82 (s, 6H), 3.54 (m, 2H), 3.28 (m, 4H), 2.74 (m, 4H), 2.16 (t, 2H), 1.45 (m, 34H), 0.88 (t, 3H). ¹³C (CDCl₃) δ 174.94, 174.08, 170.81, 162.73, 158.72, 154.29, 144.66, 144.04, 135.52, 135.28, 133.08, 132.97, 130.10, 129.99, 128.90, 128.16, 128.07, 127.90, 127.20, 113.37, 96.94, 89.29, 88.83, 86.98, 81.57, 77.23, 75.45, 74.47, 71.89, 61.01, 55.26, 39.37, 36.84, 32.22, 31.92, 29.69, 29.50, 29.36, 29.30, 29.06, 28.86, 28.47, 26.36, 25.76, 25.16, 24.70, 22.69, 14.12. MS (ES⁻) calculated for C₆₃H₈₂N₄O₁₂ 1086.6. Observed 1085.4.

Example 18**3'-O-Hexylaminopalmityl-2'-O-succinate-5'-O-DMT-N4-benzoyl-cytidine LCAA-CPG (18)**

Compound 17 (0.95 g, 0.87 mmol) and 4-dimethylaminopyridine (0.11 g, 0.87 mmol) were dissolved in CH₃CN (7.0 mL) and CH₂Cl₂ (4 mL) at room temperature in a first flask. In a second flask 2,2'-dithiobis(5-nitropyridine) (0.28

g, 0.87 mmol) was dissolved in CH₃CN (6.0 mL) and CH₂Cl₂ (2.5mL) and added to the first flask. In a third flask triphenylphosphine (0.23 g, 0.87 mmol) was dissolved in CH₃CN (7.0 mL) and then combined with the first flask. To the
5 resulting mixture was added acid washed LCAA-CPG (3.78 g, 0.44 mmol) with shaking for about 2 hours. The resulting resin was washed with CH₂Cl₂ (x3) and ether (x3). Then it was combined with Cap A (25 mL) and Cap B (25 mL) solutions from PerSeptive Biosystems GmbH, and shaken for one hour. The resin was again
10 washed with CH₂Cl₂ (x3) and ether (x3) and placed under vacuum overnight to dry. The final loading was determined to be 58 µmol/g.

Example 19

3'-O-(6-Aminohexyl-palmityl)-5'-O-DMT-uridine (19)

15 To a solution of 3'-O-(6-aminohexyl)-5'-O-DMT-uridine (1.5 g, 2.33 mmol) in CH₂Cl₂ (20 mL) at room temperature was added diisopropylamine (0.81 mL, 4.66 mmol) followed by ibuprofen pentafluorophenyl ester (compound 21, *vide infra* 1.04 g, 2.8 mmol) with stirring overnight. The mixture was
20 concentrated and the residue purified by silica gel column chromatography using EtOAc:CH₃OH (90:10; v/v) as the eluant to give 1.39 g (72%) of the title compound minus the succinyl group.

The above conjugate (1.12 mmol) succinic anhydride
25 (0.17g, 1.7 mmol), dimethylaminopyridine (0.068 g, 0.56 mmol) and triethyl amine (0.16 mL, 1.12 mmol) were dissolved in 1,2-dichloroethane (3 mL) at room temperature in a test-tube with a screw cap top. The reaction mixture was placed in a heating block at 55 °C for 2 hours and then cooled to room temperature.
30 TLC using EtOAc:MeOH (85:15; v/v) showed the absence of starting material. The mixture was diluted with 1,2-dichloroethane (30 mL) and washed three times with cold 10% aqueous citric acid (25 mL) and three times with water (25 mL).

The organic phase was dried over sodium sulfate and evaporated to give 1.2 g (quantitative yield) of the title compound as a foam.

Example 20

5 **5'-O-DMT-3'-O-ibuprofenylaminoheptyl-2'-O-succinyluridine LCAA-CPG (20)**

Compound 19 (1.02 g, 1.08 mmol) and 4-dimethylaminopyridine (0.13 g, 1.08 mmol) were dissolved in CH₃CN (9.73 mL) at room temperature in a flask. To this
10 solution was added a solution of 2,2'-dithiobis(5-nitropyridine) (0.34 g, 1.08 mmol) dissolved in acetonitrile (6.80 mL) and CH₂Cl₂ (2.90 mL) followed by a solution of triphenylphosphine (0.28 g, 1.08 mmol) dissolved in CH₃CN (9.73 mL). To this mixture was added acid washed LCAA-CPG (8.69 g,
15 with a loading of 115 mol/g) with shaking for about 2.5 hours. The resulting resin was washed with CH₃CN (x3), CH₂Cl₂ (x3), and ether (x3) to removed excess reagents. The washed resin was combined with acetic anhydride (25 mL) in THF and 1-methylimidazole (25mL) in THF (Cap A and Cap B reagents from
20 Perceptive Biosystems GmbH) with shaking for 2 hours. The resin was again washed with dichloromethane (x3) and ether (x3). Finally, it was dried overnight in a vacuum oven at room temperature under P₂O₅. The final loading was determined to be 53 mol/g.

25 A portion of the final resin (3.0 mg) was cleaved by treatment with trichloroacetic acid (25 mL, 3%) in CH₂Cl₂. The loading was determined by measuring the absorption of released trityl cation at 503 nm on a spectrophotometer (Hewlett Packard 8452A Diode Array spectrophotometer). The final derivatized
30 resin yield was 8.90 g total.

Example 21

Ibuprofenylpentafluorophenyl ester (21)

To a solution of ibuprofen (2.00 g, 9.70 mmol, Sigma) dissolved in tetrahydrofuran (20 mL) at room temperature was added 4-dimethylaminopyridine (0.24 g, 1.94 mmol) and 1, 3-dicyclohexylcarbodiimide (2.00 g, 9.70 mmol) with stirring for 20 minutes. To this mixture was added pentafluorophenol (1.78 g, 9.70 mmol) with stirring overnight. The mixture was then filtered, to remove DCU, and CH_2Cl_2 was added. The mixture was washed with water (x2), dried over magnesium sulfate, and evaporated to an oil. The oil was purified by silica gel column chromatography using ethyl acetate:hexanes (5/95, v/v) as the eluant to give 2.70 g (75%) of the title compound.

^1H NMR (CDCl_3) δ 7.21 (m, 4H), 4.05 (q, 1H), 2.47 (d, 2H), 1.86 (m, 1H), 1.63 (d, 3H), 0.90 (d, 6H). ^{13}C (CDCl_3) δ 129.65, 127.17, 45.04, 44.70, 30.18, 22.34, 18.51. MS (ES^-) calculated for $\text{C}_{19}\text{H}_{17}\text{F}_5\text{O}_2$ $[\text{M}-2\text{H}]^{2-}$ 186.1. Observed $[\text{M}-2\text{H}]^{2-}$ 183.2.

Example 22**3'-O-Hexylaminoibuprofenyl-5'-O-DMT-cytidine (22)**

To a solution of 3'-O-hexylamino-5'-O-DMT-cytidine (1.50 g, 2.33 mmol) dissolved in CH_2Cl_2 (20 mL) at room temperature was added diisopropylamine (0.81 mL, 4.66 mmol) and compound 21 (1.04 g, 2.80 mmol) with stirring overnight. The mixture was concentrated and the residue purified by silica gel column chromatography using EtOAc:MeOH (90/10; v/v) as the eluant to give 1.46 g (75%) of the title compound.

^1H NMR (CDCl_3) δ 8.00 (d, 1H), 7.39-6.80 (m, 17H), 5.90 (d, 1H), 5.80 (s, 3H), 5.48 (d, 1H), 4.29 (m, 1H), 4.18 (m, 1H), 4.02 (m, 1H), 3.78 (s, 6H), 3.62-3.31 (m, 6H), 3.13 (m, 2H), 2.43 (d, 2H), 1.83 (m, 1H), 1.48 (d, 3H), 1.40-1.18 (m, 8H), 0.88 (d, 6H). ^{13}C (CDCl_3) δ 174.54, 165.40, 158.63, 156.30, 144.43, 141.45, 140.68, 138.65, 135.52, 135.37, 130.09, 129.58, 128.13, 127.98, 127.33, 127.07, 113.24, 94.35, 91.47,

86.74, 81.32, 74.26, 70.84, 61.96, 55.22, 52.53, 46.74, 44.99, 41.06, 39.42, 30.14, 29.52, 29.34, 26.43, 25.59, 22.35, 18.50, 18.20, 12.09. MS (ES⁻) calculated for C₄₉H₆₀N₄O₈ 832.4. Observed 831.7.

5 Example 23

3'-O-Hexylaminoibuprofenyl-5'-O-DMT-N4-benzoylcytidine (23)

To a solution of compound 22 (1.45 g, 1.74 mmol) dissolved in N,N-dimethylformamide (30 mL) at room temperature was added benzoic anhydride (0.057 g, 2.53 mmol) with stirring
10 overnight. Saturated aqueous sodium bicarbonate was added and the mixture was extracted with ethyl acetate (x3). The organic phase was dried over magnesium sulfate, filtered and concentrated. The crude product was then purified by silica gel column chromatography using EtOAc:MeOH (90/10; v/v) as the
15 eluant to give 0.97 g (60%) of the title compound.

¹H NMR (CDCl₃) δ 8.75 (s, 1H), 8.41 (d, 1H), 7.91-6.84 (m, 22H), 5.99 (d, 1H), 5.35 (m, 1H), 4.37 (m, 1H), 4.26 (m, 1H), 4.08 (m, 1H), 3.81 (s, 6H), 3.62-3.36 (m, 5H), 3.15 (m, 2H), 2.44 (d, 2H), 1.84 (m, 1H), 1.50 (d, 3H), 1.45-1.16 (m,
20 8H), 0.89 (d, 6H). ¹³C (CDCl₃) δ 174.42, 162.26, 158.73, 144.06, 140.68, 138.68, 135.49, 135.26, 133.20, 130.11, 129.98, 129.61, 129.08, 128.15, 128.06, 127.52, 127.36, 127.21, 113.35, 91.81, 87.04, 81.80, 74.56, 71.01, 61.81, 56.28, 55.25, 46.80, 44.99, 39.40, 36.48, 30.18, 29.52, 29.42, 26.46, 25.63, 22.38,
25 18.50. MS (ES⁻) calculated for C₅₆H₆₄N₄O₉ 936.5. Observed 935.9.

Example 24

3'-O-Hexylaminoibuprofenyl-2'-O-succinate-5'-O-DMT-N4-benzoylcytidine (24)

Compound 23 (0.95 g, 1.01 mmol), succinic anhydride
30 (0.152 g, 1.52 mmol), dimethylaminopyridine (0.062 g, 0.50 mmol), and triethylamine (0.14 mL, 1.01 mmol) were dissolved

in 1,2-dichloroethane (4.5 mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 1 hour and then allowed to cool to room temperature. TLC using EtOAc:MeOH (90/10; v/v) 5 showed the conversion of the starting material. The mixture was diluted with CH₂Cl₂ (45 mL) washed three times with cold 10% citric acid (aqueous, 20 mL) and three times with water (20 mL). The organic-phase was dried over magnesium sulfate and evaporated to give 1.05 g (100%) of the title compound as a 10 foam.

¹H NMR (DMSO-d₆) δ 8.31 (d, 1H), 7.93 (m, 2H), 7.51-6.85 (m, 21H), 6.10 (d, 1H), 5.54 (m, 1H), 5.35 (m, 1H), 4.18 (s, 2H), 3.82 (s, 6H), 3.53 (m, 4H), 3.33-3.01 (m, 3H), 2.73 (m, 4H), 2.44 (d, 2H), 1.84 (m, 1H), 1.49 (d, 3H), 1.47-1.19 (m, 15 8H), 0.89 (d, 6H). ¹³C (CDCl₃) δ 175.31, 174.91, 170.79, 162.75, 158.72, 154.25, 144.66, 144.04, 140.76, 138.34, 135.54, 135.28, 133.06, 132.97, 130.10, 129.99, 129.64, 128.88, 128.18, 128.07, 127.92, 127.42, 127.20, 113.35, 96.94, 88.77, 86.98, 81.61, 75.47, 74.51, 71.91, 61.06, 55.24, 46.69, 44.99, 43.44, 20 39.50, 30.15, 29.71, 29.49, 29.04, 28.78, 26.24, 25.11, 22.37, 18.38. MS (ES⁻) calculated for C₆₀H₆₈N₄O₁₂ 1036.5. Observed 1035.8.

Example 25

25 3'-O-Hexylaminoibuprofenyl-2'-O-succinate-5'-O-DMT-N4-benzoylcytidine LCAA-CPG (25)

To a solution of compound 24 (1.03 g, 0.99 mmol) and 4-dimethylaminopyridine (0.12 g, 0.99 mmol) dissolved in CH₃CN (8.0 mL) at room temperature was added a solution of 2,2'-dithiobis(5-nitropyridine) (0.31 g, 0.99 mmol) dissolved in 30 CH₃CN (7.0 mL) and CH₂Cl₂ (3.0 mL) followed by a solution of triphenylphosphine (0.26 g, 0.99 mmol) dissolved in CH₃CN (8.0 mL). To the resulting mixture was added acid washed LCAA-CPG (4.31 g, 0.50 mmol) and the mixture was shaken for about 2

hours. The resulting resin was washed with CH₂Cl₂ (x3) and ether (x3). Then it was combined with Cap A (26 mL) and Cap B (26 mL) solutions from PerSeptive Biosystems GmbH, and shaken for one hour. The resin was again washed with CH₂Cl₂ (x3) and ether (x3) and dried under vacuum overnight. The final loading was determined to be 50 µmol/g.

Example 26

Synthesis of oligonucleotides incorporating compounds 14 and 20

SEQ ID NO: 1 (ISIS 22655-1 and ISIS 22656-1) and SEQ ID NO: 2 (ISIS 27700-1 and ISIS 27701-1) were synthesized on a Millipore Expedite 8901 Nucleic Acid Synthesis System.

Table I

SEQ ID NO:	ISIS #	Sequence (5'-3') ¹	U* =	Target
1	22655-1	TGC ATC CCC CAG GCC ACC AU*	cmpd 14	CD54
1	22656-1	TGC ATC CCC CAG GCC ACC AU'	cmpd 20	CD54
2	27700-1	<u>TCT</u> <u>GAG</u> <u>TAG</u> CAG AGG AGC CU*	cmpd 14	CD54
2	27701-1	<u>TCT</u> <u>GAG</u> <u>TAG</u> CAG AGG AGC CU'	cmpd 20	CD54

¹Underlined nucleosides contain 2'-O-(2-methoxyethyl) and all C's are 2'-O-(2-methoxyethyl)-5-methylcytidine.

Standard 2'-deoxy amidites (0.1M in CH₃CN, Perceptive Biosystems GmbH) were used in the synthesis of oligonucleotides having SEQ ID NO: 1 (ISIS 22655-1 and ISIS 22656-1) and oligonucleotides having SEQ ID NO: 2 (ISIS 27700-1 and ISIS 27701-1). Phosphoramidites 5'-O-DMT-2'-O-(2-methoxyethyl)-N⁶-benzoyladenine-3-O-amidite (RI Chemical), 5'-O-DMT-2'-O-(2-methoxyethyl)-N⁴-benzoyl-5-methylcytidine-3'-O-amidite (RI Chemical, Lot #E805-P-17), 5'-O-DMTr-2'-O-(2-methoxyethyl)-N²-isobutylguanosine-3'-O-amidite (RI Chemical, Lot #EMG-P-18U), and 5'-O-DMTr-2'-O-

(2-methoxyethyl)-5-methyluridine-3'-O-amidite (RI Chemical, Lot #E1050-P-10) were used in the synthesis. The 2'-O-(2-methoxyethyl) phosphoramidites were dissolved in CH₃CN (100 mg amidite/1 mL CH₃CN). Compound 14 was used as the LCAA-CPG solid support in the synthesis of SEQ ID NO: 1 (ISIS 22655-1) and SEQ ID NO: 2 (ISIS 27700-1). Compound 20 was used as the LCAA-CPG solid support in the synthesis of SEQ ID NO: 1 (ISIS 22656-1) and SEQ ID NO: 2 (ISIS 27701-1).

Each oligonucleotide was synthesized on an approximately 1 X 2 mol synthesis scale, requiring about 50 mg of derivatized LCAA-CPG for each synthesis. Deprotection of the 5'-hydroxyl groups having a DMT protecting group was performed using trichloroacetic acid (1.2 mL 3% in CH₂Cl₂) per phosphoramidite coupling followed by an CH₃CN wash. To the detritylated LCAA-CPG amidite (0.3 mL) and 1-H-tetrazole (0.6 mL, 0.49M) in CH₃CN were then delivered. The coupling time was approximately 5 minutes for standard 2'-deoxy phosphoramidites and approximately 14 minutes for novel phosphoramidites. Amidite was delivered twice per coupling. Excess amidite was washed away with CH₃CN. (2R,8aS)-(+)-(10-camphorsulfonyl) oxaziridine (0.5 mL, 36M) in CH₃CN were delivered over four minutes to oxidize the phosphodiester linkages, followed by another CH₃CN wash. Unreacted functionalities were capped with a 50:50 mixture (0.2 mL/coupling) of acetic anhydride in tetrahydrofuran (THF) and 1-methylimidazole in THF, followed by an anhydrous CH₃CN wash. Synthesis cycles (including: detritylation, amidite coupling, oxidation and capping) continued until the desired length was reached. Trityl yields were followed by the trityl monitor during the duration of each synthesis. The final DMT group was left intact.

After synthesis oligonucleotides were deprotected and cleaved from the solid support using aqueous concentrated ammonium hydroxide at 55°C for approximately 16 hours.

Oligonucleotides were then filtered from the solid support and ammonia was evaporated in a Savant AS160 Automatic Speed Vac.

The oligonucleotide crude yield was measured on a
5 Hewlett Packard 8452A Diode Array spectrophotometer at 260 nm. The crude samples were then analyzed for integrity by mass spectrometry (Hewlett Packard electrospray mass spectrometer), capillary gel electrophoresis (Beckmann P/ACE system 5000), and high performance liquid chromatography
10 (Waters 600E HPLC system with Waters 991 detector). Trityl-
on oligonucleotides were purified by HPLC (Waters) using reverse phase protocols (HPLC conditions: Waters 600E with 991 detector; Waters C₄ Delta Pak column (7.8 X 300 mm, 15 , 300 Å); solvent A = 50 mM triethylammonium acetate, pH =
15 7.0; solvent B = 100% CH₃CN; 2.5mL/minute flow rate; gradient: 5% B for first five minutes with a linear increase in B to 60% over the next 55 minutes). Appropriate HPLC fractions were collected, evaporated to completeness, detritylated in 80% acetic acid in water at room temperature
20 for approximately one hour, and then evaporated once again. To remove free trityl and excess salt, detritylated oligos were dissolved in aqueous ammonia and passed through Sephadex G-25 resin, using water as solvent. Samples were collected by a Pharmacia LKB Super Frac fraction collector.
25 The purified oligonucleotides were then analyzed for purity by CGE, MS, and HPLC (flow rate: 1.5 mL/minute, Waters Delta Pak C₄ column, 3.9 X 300 mm, 15, 300 Å). Final yields were determined by a spectrophotometer at 260 nm.

Table II

SEQ ID NO: -U*	Crude Yield (@ 260 nm)	Final Yield (@ 260 nm)	HPLC Retention Time (min) ²	Expected Mass (g/mol)	Observed Mass (g/mol)
1-14	280 ODs	104 ODs	31.18	6273.12	6270.53
1-20	324 ODs	180 ODs	37.57	6323.27	6320.99
2-14	321 ODs	137 ODs	36.08	7935.93	7929.39
2-20	303 ODs	165 ODs	36.31	7986.09	7984.52

² = HPLC conditions: Waters 600E with 991 detector
HPLC system; Waters C₄ Delta Pak column (3.9 X 300 mm, 15 ,
300 Å); solvent A = 50 mM triethylammonium acetate, pH =
7.0; solvent B = 100% CH₃CN; 1.5mL/min. flow rate; gradient:
5% B for first five minutes with a linear increase in B to
60% over the next 55 minutes. U* indicates the compound in
the specific sequence e.g. both compounds 14 and 20 were
used in each of SEQ ID NOs. 1 and 2.

Example 27

Synthesis of oligonucleotides incorporating compounds 3, 6, 9 and 12

Four oligonucleotides having SEQ ID NO: 3 (ISIS 25152-1, ISIS 25153-1, ISIS 25154-1 and ISIS 25155-1) were synthesized on a Millipore Expedite 8901 Nucleic Acid Synthesis System. The following modified amidites were used to prepare these oligonucleotides: 2'-O-methoxyethyl-thymidine (RI Chemical lot # E1050-P-10), 2'-O-methoxyethyl-5-methylcytidine (lot # S1941/RS), 2'-O-methoxyethyl-adenosine (RI Chemical, lot # EMA-P-14), and 2'-O-methoxyethylguanosine (RI Chemical, lot # EMG-P-18U). Compound 3 was used as the LCAA-CPG solid support for the synthesis of ISIS 25152-1, compound 6 for ISIS 25153-1, compound 9 for ISIS 25154-1, and compound 12 for ISIS 25155-1.

The required amounts of the amidites were placed in dried vials, dissolved in CH₃CN (modified nucleosides were prepared to give 100mg/mL), and connected to the appropriate

ports on a Millipore Expedite™ Nucleic Acid Synthesis System. solid support resin (60 mg) was used in each column for 2 X 1 μ mole scale synthesis. The synthesis was run using the standard phosphoramidite protocols utilizing (+)-
5 (2R,8aS)-10(camphorylsulfonyl)oxaziridine (CSO) for oxidation steps. The trityl reports indicated normal coupling results.

After synthesis, the oligonucleotides were deprotected with concentrated ammonium hydroxide (aq) at 55 °C for
10 approximately 16 hours, concentrated using a Savant AS160 Automatic SpeedVac, (to remove ammonia) and filtered to remove the CPG-resin. The crude samples were analyzed by MS, HPLC, and CE followed by purification on a Waters 600E HPLC system with a 991 detector (Waters C4 preparative scale
15 column) using the following solvents: A: 50 mM TEA-Ac, pH 7.0 and B: CH₃CN. The purified oligonucleotides were detritylated with 80% acetic acid at room temperature for approximately 30 minutes followed by concentrating under vacuum and drying. The oligonucleotides were dissolved in
20 concentrated ammonium hydroxide and run through a column containing Sephadex G-25 using water as the solvent and a Pharmacia LKB SuperFrac fraction collector. The resulting purified oligonucleotides were evaporated and analyzed by MS, CE and HPLC.

Table III

25

SEQ ID NO:	ISIS #	Sequence (5'-3') ¹	Modification T* =
3	25152-1	TCT GAG TAG CAG AGG AGC CT*	compound 3
30 3	25153-1	TCT GAG TAG CAG AGG AGC CT*	compound 6
3	25154-1	TCT GAG TAG CAG AGG AGC CT*	compound 9
3	25155-1	TCT GAG TAG CAG AGG AGC CT*	compound 12

All nucleotides are 2'-O-methoxyethyl (MOE) except for
T*; backbone is fully phosphodiester; and heterocycles are
35 unmodified. Are C's are 5-Me, as is the case below for

Table V.

Table IV

	SEQ ID NO. -T*	Crude Yield (@ 260 nm)	Final Yield (@ 260 nm)	HPLC/CE Retention Time (min) ²	Expected Mass (g/mol)	Observed Mass (g/mol)
5	3-3	625 ODs	288 ODs	27.7/7.30	7982.73	7982.59
	3-6	430 ODs	310 ODs	28.35/7.32	7982.73	7982.15
	3-9	480 ODs	368 ODs	27.65/7.47	7988.76	7988.41
	3-12	663 ODs	255 ODs	30.10/7.45	8002.16	8001.72

10 **Example 28****Synthesis of oligonucleotides incorporating compounds 16 (palmityl) and 21 (ibuprofenyl)**

Two oligonucleotides were synthesized having SEQ ID NO: 4 (ISIS 32361-1 and ISIS 32362-1) on a Millipore Expedite 8901 Nucleic Acid Synthesis System. Compound 16 was used as the A-CPG solid support for the synthesis of ISIS 32361-1 and also a palmityl TC dimer. Compound 21 was used as the LCAA-CPG solid support for the synthesis of ISIS 32362-1 and an ibuprofenyl TC dimer. The following modified amidites were used in the above sequences: 5'-DMT-2'-O-methoxyethyl-5-methyluridine beta-cyanoethylphosphoramidite (PrOligo, Lot No. S 3044), 2'-O-(2-methoxyethyl)-5-Me-C Bz amidite (BSR-1026-89), 2'-O-MOE A phosphoramidite (Pharmacia Biotech, Lot No. 311119), and 2'-O-(2-methoxyethyl)-(iBu)G amidite (BSR-1026-84).

The required amounts of the amidites were placed in dried vials, dissolved in CH₃CN (modified nucleosides prepared to be 100 mg/mL), and connected to the appropriate ports on a Millipore Expedite™ Nucleic Acid Synthesis System. Solid support resin (60 mg) was used in each column for 2 X 1 μmol scale synthesis. The synthesis was run using standard phosphoramidite protocols utilizing CSO for

oxidation steps. The trityl reports indicated normal coupling results. After synthesis, the oligonucleotides were deprotected with concentrated ammonium hydroxide (aq) at 55 °C for approximately 16 hours. Then they were
 5 evaporated, using a Savant AS160 Automatic SpeedVac, (to remove ammonia) and filtered to remove the CPG-resin.

The crude samples were analyzed, purified and deprotected as illustrated above in Example 27. The dried oligonucleotides were dissolved in concentrated ammonium
 10 hydroxide and run through a column containing Sephadex G-25 with water used as eluent. The dimers were each further purified using a Dowex and then a Chelex column for NMR studies. The resulting purified oligonucleotides were evaporated and analyzed by MS, CE (MDQ) and HPLC.

15

Table V

SEQ ID NO:	ISIS #	Sequence (5'-3') ¹	Modification C* =
4	32361-1	TCT GAG TAG CAG AGG AGC TC*	compound 16
20 4	32362-1	TCT GAG TAG CAG AGG AGC TC*	compound 21
dimer		TC* (16)	compound 16
dimer		TC* (21)	compound 21

All nucleotides are 2'-O-methoxyethyl (MOE) except for C*; backbone is fully phosphodiester; and heterocycles are
 25 unmodified except all C's are 5-methylcytidine.

TABLE VI

SEQ ID NO: -C*	Crude Yield (@ 260 nm)	Final Yield (@ 260 nm)	HPLC/CE Retention Time (min) ²	Expected Mass (g/mol)	Observed Mass (g/mol)
30 4-16	498 ODs	95 ODs	45.71/4.325	7987.59	7984.86
4-21	649 ODs	122 ODs	32.45/4.500	7937.48	7932.63
dimer-16	33 ODs	22 ODs	n/a	M ²⁺ - 593.6	M ²⁺ - 592.7

dimer-21	30 ODs	20 ODs	n/a	M ²⁻ 568.5	M ²⁻ 567.5
----------	--------	--------	-----	-----------------------	-----------------------

Example 29**Synthesis of oligonucleotides incorporating compounds 18****5 (palmityl) and 20 (ibuprofenyl)**

Following the procedures illustrated in the examples above compounds 18 and 20 were incorporated into oligonucleotides SEQ ID NO: 4 as illustrated in Table VII. Compound 4-18 is ISIS 32361-1 and compound 4-20 is ISIS
 10 32362-1.

Table VII

SEQ ID NO:	Sequence ⁴	Back-bone	ODs (@260 nm)
4-18	TC ^{5Me} T GAG TAG C ^{5Me} AG AGG AGC ^{5Me} TC*	P=O	30
15 4-20	TC ^{5Me} T GAG TAG C ^{5Me} AG AGG AGC ^{5Me} TC**	P=O	30

⁴All nucleosides contain the 2'-O-Methoxyethyl group (except for the 3'-terminal C). All C's are 5-methyl-C's. C* = 3'-O-Palmityl-aminoheptyl-cytidine. C** = 3'-O-Ibuprofenyl-aminoheptyl-cytidine.

20 Example 30**Synthesis of oligonucleotides incorporating compound (26)**

Compound 26 was incorporated into SEQ ID NO: 4 (ISIS 29782-1) as the 3'-terminal nucleoside. The synthesis was performed on a Millipore Expedite 8901 nucleic acid
 25 Synthesizer. The incorporation of compound 26 into an oligonucleotide allows the conjugation at the 3'-end of the oligonucleotide via the 2'-aminopropyl group.

Table VIII

SEQ ID NO:	Sequence (5'-3') ⁵	C* =	Target
30 -C*			

4-26	TCT GAG TAG CAG AGG AGC TC*	compound 26	CD54
------	-----------------------------	----------------	------

⁵All nucleosides contain 2'-O-(2-methoxyethyl) except C*. All C's are 2'-O-(2-methoxyethyl)-5-methylcytidine.

Nucleosides were purchased from commercial sources: 5'-O-DMT-2'-O-(2-methoxyethyl)-N⁶-benzoyladenine-3'-O-amidite (RI Chemical, Lot #EMA-P-09); 5'-O-DMT-2'-O-(2-methoxyethyl)-N⁴-benzoyl-5-methylcytidine-3'-O-amidite (RI Chemical, Lot #E805-P-17); 5'-O-DMT-2'-O-(2-methoxyethyl)-N²-isobutylguanosine-3'-O-amidite (Pharmacia Biotech 27-0022-42), and 5'-O-DMT-2'-O-(2-methoxyethyl)-5-methyluridine-3'-O-amidite (Perceptive Biosystems). The 2'-O-(2-methoxyethyl) phosphoramidites were dissolved in CH₃CN (100 mg amidite/1 mL CH₃CN). Compound 26 was used as the LCAA-CPG solid support in the synthesis which effected its incorporation at the 3' end of the oligonucleotide.

The oligonucleotide was synthesized on an approximately 2 X 20 μ mol synthesis scale, requiring approximately 333 mg of derivatized LCAA-CPG each. The DMT protecting groups on the solid support were removed with tRI Chemicalchloroacetic acid (10.6 mL, 3%) in CH₂Cl₂ per coupling followed by an CH₃CN wash. To the detritylated LCAA-CPG, amidite (1.20 mL) and 1-H-tetrazole (1.80 mL, 0.49M) in CH₃CN were then delivered (total coupling time of approximately 24 minutes for novel amidites.) The amidite reagent was delivered four times per coupling. Excess amidite was washed away with CH₃CN. (2R, 8aS)-(+)-(10-camphorsulfonyl) oxaziridine (2.40 mL 36.4M) in anhydrous CH₃CN was delivered over four minutes to oxidize the phosphodiester linkages, followed by another anhydrous CH₃CN wash. Unreacted functionalities were capped with a 50:50 mixture (1.40 mL/coupling) of acetic anhydride in tetrahydrofuran (THF) and 1-methylimidazole in THF, followed

by an anhydrous CH_3CN wash. Trityl yields were followed by the trityl monitor during the duration of a synthesis. The final DMT group was left intact.

Following the synthesis, the oligonucleotides were
5 deprotected and removed from the solid support with concentrated ammonium hydroxide (aq) and methylamine (Aldrich Chemicals, 10%, 40 wt. % solution in water) at 55 °C for approximately 16 hours. They were then filtered from the solid support and ammonia was evaporated in a
10 Savant AS160 Automatic Speed Vac.

Oligonucleotide crude yield was measured on a Hewlett Packard 8452A Diode Array spectrophotometer at 260 nm. The crude samples were then analyzed for integrity by mass spectrometry (Hewlett Packard electrospray mass
15 spectrometer), capillary gel electrophoresis (Beckmann P/ACE system 5000), and high performance liquid chromatography (Waters 600E HPLC system with Waters 991 detector). Trityl-on oligonucleotides were purified on the Waters HPLC system by reverse phase as illustrated above. (HPLC conditions:
20 waters C_4 Delta Pak column (25 X 100 mm, 15 , 300 Å); solvent A = 50 mM triethylammonium acetate, pH = 7.0; solvent B = 100% CH_3CN ; 5.0mL/min. flow rate; gradient: 5% B for first five minutes with a linear increase in B to 60% over the next 55 minutes.) Appropriate HPLC fractions were
25 collected, evaporated to completeness, detritylated in 80% acetic acid in water at room temperature for approximately one hour, and then evaporated once again. To remove free trityl and excess salt, detritylated oligos were dissolved in aqueous ammonia and passed through Sephadex G-25 resin,
30 using water as solvent. Samples were collected by a Pharmacia LKB Super Frac fraction collector. The purified oligonucleotides were then analyzed for purity by CGE, MS, and HPLC (flow rate: 1.5 mL/min., Waters Delta Pak C_4 column, 3.9 X 300 mm, 15, 300 Å). Final yields were

determined by a spectrophotometer at 260 nm.

Table IX

SEQ ID NO: -C*	Crude Yield (@ 260 nm)	Final Yield (@ 260 nm)	HPLC Retention Time (min)	Expected Mass (g/mol)	Observed Mass (g/mol)
4-26	6902 ODs	3100 ODs	22.19	7705.25	7704.32

Example 31**Ligand conjugation to an oligonucleotide containing an****aminopropyl linker**

Oligonucleotide having SEQ ID NO: 4 and further having compound 26 attached as the 3'-nucleoside (4-26) was used as a substrate for post synthetic conjugation of functional groups. Four different functional groups (PEG₂₀₀₀, PEG₅₀₀₀, Biotin, and Pyrene) were conjugated and the respective oligonucleotides were purified. The groups are attached at the 3' end of the oligonucleotide via a 2-O-aminohexyl linking group.

Table X

SEQ ID NO:	Sequence (5'-3')	Modification	Target
4-26	TCT GAG TAG CAG AGG AGC TC'	C' = 2'-O-PEG ₂₀₀₀ - aminopropylcytidine	CD54
4-26	TCT GAG TAG CAG AGG AGC TC'	C' = 2'-O-PEG ₅₀₀₀ - aminopropylcytidine	CD54
4-26	TCT GAG TAG CAG AGG AGC TC'	C' = 2'-O- biotinylaminopropylcytidine	CD54
4-26	TCT GAG TAG CAG AGG AGC TC'	C' = 2'-O-pyrenylpropyl carbonylaminopropylcytidine	CD54

All nucleotides are 2'-O-MOE modified except C*.

All C's are 2'-O-(2-methoxyethyl)-5-methylcytidine.

A) Procedure for PEG₂₀₀₀, ISIS 30130-1

ISIS 29782-1 (100 ODs) contained in a closed-capped 13 X 100 mm pyrex test tube was dried down in a speed vac overnight. After drying, 200 mg PEG₂₀₀₀ and sodium bicarbonate (400 μ L, 0.2M) were added to the oligonucleotide with shaking overnight. The reaction mixture was dissolved in water (3 mL) and purified by HPLC (HPLC conditions: Waters C₄ Delta Pak column (7.8 X 300 mm, 15 , 300 Å); solvent A = 50 mM triethylammonium acetate, pH = 7.0; solvent B = 100% CH₃CN; flow rate 2.5 mL/minute; gradient: 5% B for first five minutes with a linear increase in B to 60% over the next 55 minutes. The fractions of interest were collected and evaporated. To remove salt and free PEG₂₀₀₀ the oligonucleotide was passed through Sephadex G-25 resin and further purified by HPLC (conditions: solvent A = 50 mM triethylammonium acetate, pH = 7.0; solvent B = 100% CH₃CN; solvent C = H₂O; flow rate 2.5 mL/minute; gradient: 100% A for first 10 minutes with a linear increase in C to 100% over the next 5 minutes, remaining constant for the next 60 minutes, followed by a linear increase in B to 100% for 20 minutes.) ISIS 30130-1 was analyzed for purity by Mass Spec, HPLC, and CGE. Final yield was determined by spectrophotometer at 260 nm.

B) Procedure for PEG₅₀₀₀, ISIS 30131-1

ISIS 29782-1 (100 ODs) contained in a closed-capped 13 X 100 mm pyrex test tube was dried down in a speed vac overnight. After drying, 150 mg PEG₅₀₀₀ and sodium bicarbonate (350 μ L, 0.2M) was added with shaking overnight. The reaction mixture was dissolved in water (3 mL) and purified by HPLC as illustrated above. The final oligonucleotide was analyzed for purity by Mass Spec, HPLC,

and CGE. Final yield was determined by spectrophotometer at 260 nm.

C) Procedure for biotin, ISIS 30132-1

ISIS 29782-1 (100 ODs) contained in a closed-capped 13 X 5 100 mm pyrex test tube was dried down in a speed vac overnight. After drying, 20 mg (+)-biotin N-succinimidyl ester (Fluka 14405) and sodium bicarbonate (200 μ L 0.2M) was added to the oligonucleotide with shaking overnight. The mixture was shaken overnight. The reaction mixture was 10 dissolved in water (3 mL) and purified by HPLC as illustrated above. The final oligonucleotide was analyzed for purity by Mass Spec, HPLC, and CGE. Final yield was determined by spectrophotometer at 260 nm.

D) Procedure for pyrene, ISIS 30133-1

ISIS 29782-1 (100 ODs) contained in a closed-capped 13 X 100 mm pyrex test tube was dried down in a speed vac overnight. After drying, 20 mg succinimidyl-1-pyrene butyrate (Molecular Probes, Lot #2721-3) and sodium 20 bicarbonate (200 μ L, 0.2M) was added to the oligonucleotide. The reaction mixture was dissolved in water (3 mL) and purified by HPLC as illustrated above. The final oligonucleotide was analyzed for purity by Mass Spec, HPLC, and CGE. Final yield was determined by spectrophotometer at 25 260 nm.

Table XI

ISIS #	Starting Yield	Final Yield	HPLC Retention Time (min) ²	Expected Mass (g/mol)	Observed Mass (g/mol)
	(@ 260 nm)	(@ 260 nm)			
30 30130-1	100 ODs	28 ODs	32.61	N/A	N/A
30131-1	100 ODs	30 ODs	39.14	N/A	N/A

30132-1	100 ODs	26 ODs	24.17	7932.56	7930.49
30133-1	100 ODs	25 ODs	22.43	7976.59	7976.63

Example 32**5 3'-O-Hexylaminobenzylpenicillinyl-5'-O-DMT-5-methyluridine (22)**

Benzylpenicillin potassium salt (0.56 g, 1.52 mmol, Fluka) was suspended in DMF (6 mL) at room temperature under an atmosphere of argon. 4-Methylmorpholine (0.33 mL, 3.04 mmol) and TBTU (0.49 g, 1.52 mmol) were added and the suspension became a clear orange solution. 3'-O-Hexylamino-5'-O-DMT-5-methyluridine (1.0 g, 1.52 mmol) was added and the mixture with stirring overnight. The mixture was evaporated under high vacuum to give the title compound.

15 Example 33**3'-O-Hexylaminophenoxymethylpenicillinyl-5'-O-DMT-5-methyluridine (23)**

Phenoxymethylpenicillinic acid (1.06 g, 3.03 mmol, Sigma) was dissolved in DMF (10 mL) at room temperature under an atmosphere of argon. 4-Methylmorpholine (0.67 mL, 6.06 mmol) and TBTU (0.97 g, 3.03 mmol) were added followed by 3'-O-hexylamino-5'-O-DMT-5-methyluridine (2.0 g, 3.03 mmol). The mixture was stirred overnight and then evaporated. The material was purified by silica gel column chromatography using ethyl acetate:triethylamine (100/1, v/v) as the eluant to give 0.496 g of the title compound.

Example 34**Succinimidylphenoxymethylpenicillin (24)**

Phenoxymethylpenicillinic acid (1.00g, 2.85 mmol, Sigma) was suspended in CH₂Cl₂ (10 mL) at room temperature

under an atmosphere of argon. Dimethylaminopyridine (DMAP) (0.070 g, 0.57 mmol) was added and the suspension dissolved into a clear solution. 1,3-Dicyclohexylcarbodiimide (0.59 g, 2.85 mmol) was added with stirring for about 30 minutes followed by addition of N-hydroxysuccinimide (0.33 g, 2.85 mmol). The suspension stirred for about 3 hours and filtered to remove DCU. The filter cake was washed with CH_2Cl_2 and the combined organic phase was washed twice with water (to remove DMAP). The organic-phase was then dried over sodium sulfate and evaporated to a brown foam to give 1.26 g (98%) of the title compound. MS (ES^+) calculated for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_7\text{S}$ 447.1. Observed MH^+ 449.1 (minor peak) and MH^{2+} 224.9 (major peak).

15 Example 35

Preparation of phenoxyethylpenicillinyl conjugated oligonucleotide SEQ ID NO: 5

5'-Hexanolamine-phosphodiester-TGC ATC CCC CAG GCC ACC AT, SEQ ID NO: 5, (ISIS 3082) was prepared following standard methods and techniques using an automated DNA synthesizer. At the last step in synthesis 5'-amino-modifier C6 phosphoramidite (Glen Research, Sterling, VA) was used to introduce the aminohexylphosphodiester attached to the 5'-end of the oligomer. All internucleotide linkages were P=O (phosphodiester) linkages and they were introduced by CSO oxidation protocol. The final oligonucleotide was deprotected and HPLC purified to give the 5'-aminohexyl phosphodiester linked oligonucleotide.

The aminohexyl linked oligonucleotide was dried to a white powder and dissolved in sodium bicarbonate (200 μL , 0.1 M, aq) at room temperature. Compound 24 (25 mg, 0.06 mmol) in DMF (200 μL) was added with vortexing and the mixture was kept overnight at room temperature. The mixture was run through a sephadex G-25 column using water as the

solvent. The collected fractions were filtered through a syringe disk filter and purified by prep-HPLC using a C-4 column as illustrated in the previous examples. The collected fractions were concentrated and dried to give the
5 title oligonucleotide.

Example 36

Preparation of phenoxymethylpenicillinyl conjugated full 2'-O-MOE oligonucleotide SEQ ID NO: 3

5'-Hexanolamine-phosphodiester-TC^{5MT} GAG TAG C^{5MAG} AGG
10 AGC^{5M} C^{5MT}, SEQ ID NO: 3, (ISIS 11158) was prepared following standard methods and techniques using an automated DNA synthesizer. At the last step in the synthesis 5'-amino-modifier C6 phosphoramidite (Glen Research, Sterling, VA) was used to introduce the aminohexylphosphodiester attached
15 to the 5'-end of the oligomer. All internucleotide linkages were P=O (phosphodiester) linkages introduced by CSO oxidation protocol. The final oligonucleotide was deprotected and HPLC purified to give the 5'-aminohexyl phosphodiester linked oligonucleotide.

20 The aminohexyl linked oligonucleotide (50 OD's) was dissolved in sodium bicarbonate (200 μ l, 0.1 M, aq) at room temperature. Compound 24 (25 mg, 0.06 mmol) in DMF (100 μ L) was added, the resulting suspension was vortexed and allowed to stand overnight at room temperature. The mixture was run
25 through a sephadex G-25 column using water as the solvent. The collected fractions were filtered through a syringe disk filter and purified through a prep-HPLC using a C-4 column as illustrated above. These collected fractions were then evaporated to give the title oligonucleotide.

30

Example 37

Preparation of 3'-O-hexylaminoaspirinyl-5'-O-DMT-5-

methyluridine 25

Acetyl salicylic acid (aspirin) (0.55 g, 3.03 mmol) was dissolved in DMF (10 mL) at room temperature under an atmosphere of argon. 4-Methylmorpholine (0.67 mL, 6.06 mmol) and TBTU (0.97 g, 3.03 mmol) were added followed by 3'-O-hexylamino-5'-O-DMT-5-methyluridine (2.00 g, 3.03 mmol). The mixture was stirred overnight and concentrated. The crude material was purified by silica gel column chromatography using ethyl acetate:hexanes:triethylamine (75/25/1, v/v/v) as the eluant to give 1.36 g (55%) of the title compound as a clear oil. MS (ES⁺) calculated for C₄₆H₅₁N₃O₁₁ 821.4; observed MH⁺ + Na 844.4.

Example 38**3'-O-Hexylaminoaspirinyl-2'-O-succinyl-5'-O-DMT-5-****15 methyluridine 26**

Compound 25 (1.31 g, 1.59 mmol) was dissolved in 1,2-dichloroethane (4 mL) at room temperature under an atmosphere of argon. Triethylamine (0.22 mL, 1.59 mmol), DMAP (0.097g, 0.80 mmol), and succinic anhydride (0.239 g, 2.38 mmol) were added and the mixture was placed in a 50°C heating block overnight to give following purification the title compound.

Example 40**Succinimidylaspirin 27**

25 Acetylsalicylic acid (1.00g, 5.55 mmol) and DMAP (0.136g, 1.11 mmol) were dissolved in CH₂Cl₂ (10 mL) at room temperature under an atmosphere of argon. DCC (1.145 g, 5.55 mmol) was added, the mixture was stirred for about 5 minutes and N-hydroxysuccinimide (0.639 g, 5.55 mmol) was added. The mixture stirred for 4 hours, filtered, CH₂Cl₂ was added and the mixture was washed twice with water. The organic phase was dried over sodium sulfate, concentrated

and dried to give 1.58 g (100%) of the title compound.

Example 41

Preparation of aspirinyl conjugated oligonucleotide SEQ ID NO: 3

5 5'-Hexanolamine-phosphodiester-TC^{5M}T GAG TAG C^{5M}AG AGG
AGC^{5M} C^{5M}T, SEQ ID NO: 3, prepared as per Example 36, (ISIS
11158) (100 OD's) (dried to a white powder) was dissolved in
sodium bicarbonate (0.1 M, 200 µL, aq) at room temperature.
Compound 38 (25 mg, 0.06 mmol) in DMF (400 µL) was added and
10 the resulting suspension was vortexed and then shaken
overnight at room temperature. The resulting material was
run through a sephadex G-25 column using water as the
eluent. The collected fractions were filtered through a
syringe disk filter and purified through a prep-HPLC C-4
15 column as illustrated above to give after concentration and
drying the title oligonucleotide.

Example 42

Binding affinity of oligonucleotides to human serum albumin (HSA)


20 Binding curves:

The 5' end of ISIS-27700 was end labeled with ³²P using
T4 polynucleotide kinase and standard procedures.
Unincorporated label was removed using a G25 column and was
confirmed by polyacrylamide gel electrophoresis. A fixed
25 concentration of labeled oligonucleotide (50 nM) was
incubated with increasing concentrations of human serum
albumin (Fraction V, essentially Fatty Acid Free,
essentially globulin free, Sigma Chemical Company, St.
Louis, MO) and incubated at 25° for one hour in PBS plus 0.1
30 mM EDTA and 0.005% Tween 80. Experiments with longer
incubation times demonstrated that full equilibrium was

achieved in less than one hour.

Albumin-oligonucleotide mixtures were placed on membranes (Ultrafree-MC 30000, Millipore) and spun very gently at 3000 rpm (725xg) for 3-6 minutes until ~20% of the volume had passed through the filter. Aliquots of the initial mix (before filtration) and the filtrates were counted in a scintillation counter. After appropriate correction for background, the concentration of free and bound oligonucleotide was calculated.

10 The low concentration of oligonucleotide, relative to albumin, allows for detection of binding to only the tightest binding site on the albumin. Thus, the fraction of bound oligonucleotide was plotted versus the total albumin concentration and data were fit to a two state model:



where O is unbound oligonucleotide, A is unbound albumin, (OA) is the oligonucleotide-albumin complex and K_A is the equilibrium association constant.

Capacity curves:

20 Capacity curves were measured using a technique similar to that used for the binding curves except a fixed concentration of albumin (50 μ M) was employed and the concentration of labeled oligonucleotide was varied. The average number of moles of oligonucleotide bound per mole of protein, n_L , was plotted versus free oligonucleotide concentration and fit to a model with two classes of binding sites, each with n_i binding sites per protein and association constant, K_i .

Results:

30 Oligonucleotides tested are listed in Table XII. A

comparison was made between an unmodified deoxy diester oligonucleotide (8651) and its 3' ibuprofen conjugate (22655) and a uniformly 2'-O-methoxy-ethyl modified phosphodiester oligonucleotide (11158) and its 3' ibuprofen conjugate (27700). As seen in Figure 1 and Table XIII, binding of the unconjugated controls was very weak ($K_D > 200 \mu\text{M}$). Addition of the ibuprofen conjugate increased the affinity substantially. Binding of the phosphodiester conjugates was comparable to that of phosphorothioate DNA oligonucleotides which are among the tightest binding of all modified oligonucleotides (data not shown). The capacity of HSA for the ibuprofen conjugate was also measured. Binding ratios of 0.75:1 (oligonucleotide: albumin) were achieved for the conjugate. This contrasts to unconjugated oligonucleotides where maximum capacity observed was 0.2:1.

Conclusion:

Phosphodiester oligonucleotides (both 2'-deoxy and 2'-O-methoxyethyl) bound to HSA with weak affinity ($K_D > 200 \mu\text{M}$). Phosphorothioate oligonucleotides, in contrast, had greater affinities (K_D 3-30 μM). Addition of an ibuprofen conjugate to the 3' end of a phosphodiester oligonucleotide increased the affinity into the range typical for phosphorothioate oligonucleotides. It was seen that the capacity of HSA for the ibuprofen conjugate was much greater than that for unconjugated oligonucleotides.

Table XII

SEQ ID	ISIS #	Sequence	Chemistry	Conjugate
NO:				
3	27700	TC*T GAG TAG C*AG full MOE PO AGG AGC* C*T		3'-ibuprofen
4	11158	TC*T GAG TAG C*AG full MOE PO AGG AGC* TC		3'-OH

5	4	3067	TCT GAG TAG CAG AGG AGC TC	full deoxy PS	3'-OH
	5	22655	TGC ATC CCC CAG GCC ACC AT	full deoxy PO	3'-ibuprofen
	5	8651	TGC ATC CCC CAG GCC ACC AT	full deoxy PO	3'-OH
	5	3082	TGC ATC CCC CAG GCC ACC AT	full deoxy PS	3'-OH
			C* = 5-methyl cytosine.		

10

Table XIII

Equilibrium dissociation constants for modified
oligonucleotides binding to HSA.

	SEQ ID NO:	ISIS #	K _d (μM)
15	3	27700	8
	4	11158	>400
	4	3067	7
	5	22655	12
	5	8651	>200
	5	3082	4

20 *Equilibrium constants were obtained from data in Figure
1 as described in the text.

Example 43**2'-O-Hexylaminopalmitoyl-5'-O-DMT-adenosine 28**

2'-O-Hexylamino-5'-O-DMT-adenosine (3.00 g, 4.49 mmol)
25 was dissolved in dichloromethane (60 mL, anh) at room
temperature. Diisopropylamine (1.56 mL, 8.98 mmol) and
palmitic acid pentafluorophenyl-ester (2.28 g, 5.39 mmol)
were added and the mixture was stirred overnight and
evaporated. The crude material was purified on a silica
30 column (250 mL) using EtOAc-MeOH (95:5) as the eluent to
give ~4.07 g (~100%) of the title compound.

Example 44

2'-O-Hexylaminopalmityl-5'-O-DMT-N⁶-Benzoyladenosine 29

Compound 28 (4.00 g, 4.41 mmol) was dissolved in anhydrous pyridine (50 mL) at room temperature under argon. The solution was cooled to ice temperature and 5 chlorotrimethylsilane (1.40 mL, 11.02 mmol) was added. The mixture stirred at ice temperature for ~30 minutes when benzoyl chloride (1.54 mL, 13.23 mmol) was added. Then it was allowed to warm to room temperature and stirred overnight. The mixture was cooled to ice temperature again 10 and cold water (10 mL) was added. It was stirred for 15 minutes, then cold concentrated ammonium hydroxide (10 mL) was added. The mixture was allowed to warm to room temperature and stirred for 30 minutes, after which it was evaporated. Water (25 mL) was added and the mixture was 15 extracted with ethyl acetate (x3). The organic phase was dried over anhydrous sodium sulfate and evaporated. A 300 mL silica column was run using ethyl acetate-hexanes (50:50) as the solvent to yield 1.83 g (41%) of the title compound.

Example 45**20 2'-O-Hexylaminopalmityl-3'-O-succinate-5'-O-DMT-N⁶-Benzoyladenosine 30**

Compound 29 (1.19 g, 1.18 mmol), succinic anhydride (0.22 g, 1.77 mmol), dimethylaminopyridine (0.09 g, 0.59 mmol), and triethylamine (0.21 mL, 1.18 mmol) were dissolved 25 in 7 mL of 1,2-dichloroethane at room temperature. The reaction mixture (in a test-tube with a screw cap top) was placed in a heating block at 55 °C for 1 hour and then allowed to cool to room temperature overnight. TLC (EtOAc-hexanes, 90:10) showed the absence of starting material. 30 Dichloromethane (70 mL) was added and the mixture was washed three times with 30 mL portions of cold 10% aqueous citric acid followed by three washes with 30 mL portions of water.

The organic-phase was dried over anhydrous sodium sulfate and evaporated to afford 1.26 g (97%) of the title compound as a foam.

Example 46

5 2'-O-Hexylaminopalmityl-3'-O-succinate-5'-O-DMT-N⁶-benzoyl-adenosine LCAA-CPG 31

Compound 30 (1.24 g, 1.12 mmol) and 4-dimethylamino-pyridine (0.14 g, 1.12 mmol) were dissolved in acetonitrile (7.0 mL, anh) at room temperature. In another flask, 2,2'-
10 dithiobis-5-nitropyridine (0.35 g, 1.12 mmol) was dissolved in anhydrous acetonitrile (4.0 mL) and anhydrous dichloromethane (4.0 mL). This solution was then added to the first flask. In a third flask, triphenylphosphine (0.29
15 g, 1.12 mmol) was dissolved in acetonitrile (6.0 mL) and then combined with the first flask. Finally, acid-washed LCAA-CPG (4.86 g, 0.56 mmol) was added and the mixture was shaken for ~2 hours. The resulting resin was washed three times with dichloromethane and ether. Then it was combined with Cap A (21 mL) and Cap B (21 mL, solutions from
20 PerSeptive Biosystems GmbH) and shaken for an additional hour. The resin was then washed again three times with dichloromethane and ether and placed under vacuum overnight to dry. The final loading was determined to be 48 umol/g.

Example 47

25 2'-O-Hexylaminoibuprofenyl-5'-O-DMT-adenosine 32

2'-O-Hexylamino-5'-O-DMT-adenosine (3.00 g, 4.49 mmol, RI Chemical Company) was dissolved in anhydrous dichloromethane (40 mL) at room temperature. Diisopropylamine (1.56 mL, 8.98 mmol) and ibuprofen-
30 pentafluorophenylester (2.01 g, 5.39 mmol, Example 21) were added and the mixture stirred overnight and evaporated. The

crude material was evaporated and purified over a 250 mL silica column using EtOAc-MeOH (95:5) as the solvent to give 2.89 g (75%) of the title compound.

Example 48

5 2'-O-Hexylaminoibuprofenyl-5'-O-DMT-N⁶-Benzoyladenosine 33

Compound 32 (2.87 g, 3.35 mmol) was dissolved in anhydrous pyridine (50 mL) at room temperature under Ar(g). The solution was cooled to ice temperature and chlorotrimethylsilane (1.06 mL, 8.38 mmol) was added. The mixture stirred at ice temperature for ~30 min. and then benzoyl chloride (1.17 mL, 10.05 mmol) was added. The mixture was allowed to warm to room temperature and stirred overnight. The mixture was cooled to ice temperature again and cold water (10 mL) was added with stirring for 15 minutes. Cold concentrated ammonium hydroxide (10 mL) was then added. The mixture was allowed to warm to room temperature and stirred for 30 minutes and evaporated. Water (25 mL) was added and the mixture was extracted with ethyl acetate (x3). The organic phase was dried over sodium sulfate and evaporated. The resulting material was purified using a 200 mL silica column with ethyl acetate-hexanes (90:10) as the eluant to give 2.50 g (78%) of the title compound.

Example 49

25 2'-O-Hexylaminoibuprofenyl-3'-O-succinate-5'-O-DMT-N⁶-benzoyladenosine 34

Compound 33 (2.00 g, 2.08 mmol), succinic anhydride (0.312 g, 3.12 mmol), dimethylaminopyridine (0.127 g, 1.04 mmol), and triethylamine (0.29 mL, 2.08 mmol) were dissolved in 1,2-dichloroethane (9 mL) at room temperature. The reaction mixture (in a test-tube with a screw cap top) was

placed in a heating block at 55 °C for 1 hour and then allowed to cool to room temperature overnight. TLC (EtOAc-MeOH, 90:10) showed the absence of starting material.

Dichloromethane (90 mL) was added and the mixture was washed 5 three times with 40 mL portions of cold 10% aqueous citric acid followed by three washes with 40 mL portions of water. The organic-phase was dried over anhydrous sodium sulfate and evaporated to a foam to give 1.86 g (84%) of the title compound.

10 Example 50

2'-O-Hexylaminoibuprofenyl-3'-O-succinate-5'-O-DMT-N⁶-benzoyladenine LCAA-CPG 35

Compound 34 (1.80 g, 1.70 mmol) and 4-dimethylamino-
pyridine (0.21 g, 1.70 mmol) were dissolved in anhydrous
15 acetonitrile (10.0 mL) at room temperature. In another
flask, 2,2'-dithiobis(5-nitropyridine) (0.53 g, 1.70 mmol)
was dissolved in anhydrous acetonitrile (7.0 mL) and
anhydrous dichloromethane (6.0 mL). This solution was then
added to the first flask. In a third flask,
20 triphenylphosphine (0.45 g, 1.70 mmol) was dissolved in
acetonitrile (8.0 mL) and then combined with the first
flask. Finally, acid-washed LCAA-CPG (7.38 g, 0.85 mmol)
was added and the mixture was shaken for ~2 hours. The
resulting resin was washed three times with dichloromethane
25 and ether. Then it was combined with Cap A (30 mL) and Cap
B (30 mL) solutions from PerSeptive Biosystems GmbH and
shaken for an additional hour. The resin was then washed
again three times with dichloromethane and ether and placed
under vacuum overnight to dry. The final loading was
30 determined to be 50 $\mu\text{mol/g}$.

Example 51

Procdeure for SEQ ID NO's. 6 and 7

The following modified amidites were used in the above sequences: 2'-O-methoxyethyl-thymidine (RIC, Inc., lot # E1050-P-10), 2'-O-methoxyethyl-5-methylcytidine (lot # 5 S1941/RS), 2'-O-methoxyethyl-adenosine (lot # EMA-P-14 RIC), and 2'-O-methoxyethyl-guanosine (lot # EMG-P-18U RIC). Compound 35 (which is attached to CPG) was used as the LCAA-CPG solid support for the synthesis of Isis #111494-1 and 111496-1. MDC-1395-94 (compound 31) was used as the LCAA-10 CPG solid support for the synthesis of Isis #111495-1 and 111497-1.

The required amounts of the amidites (1 M solutions of unmodified nucleosides and 100 mg/mL of modified nucleosides) were placed in dried vials, dissolved in 15 acetonitrile, and connected to the appropriate ports on a Millipore Expedite™ Nucleic Acid Synthesis System (ISIS 4). 60mg of solid support resin was used in each column for 2X1 umole scale synthesis. The synthesis was run using the IBP-PS(1umole) double coupling protocol for phosphorothioate 20 backbones. The trityl reports indicated normal coupling results.

After synthesis, the oligonucleotides were deprotected with conc. ammonium hydroxide(aq) at 55 °C for approximately 16 hrs. Then they were evaporated, using a Savant AS160 25 Automatic SpeedVac, (to remove ammonia) and filtered to remove the CPG-resin.

The crude samples were analyzed by MS, HPLC, and CE. Then they were purified on a Waters 600E HPLC system with a 991 detector using a Waters C18 Prep. scale column (C18 30 Prep.) and the following solvents: A: 0.1 M aqueous ammonium acetate and B: acetonitrile utilizing the "C18PREP" method.

After purification the oligos were evaporated to dryness and then detritylated with 80% acetic acid at room temperature for approximately 30 minutes and again

evaporated. The oligonucleotides were desalted by dissolving them in water with concentrated ammonium hydroxide and running them through a C18 Prep. column using water as the solvent. The oligonucleotides were then washed
 5 from the column with acetonitrile. The resulting purified oligonucleotides were evaporated and analyzed by MS, CE and HPLC.

Table XIV

10	SEQ ID	ISIS #	Backbone	Sequence ¹
	NO.			
	6	111494	P=S	GTT C ^{5Me} TC ^{5Me} GC ^{5Me} T GGT GAG TTT C ^{5Me} A ^{IBU}
	6	111495	P=S	GTT C ^{5Me} TC ^{5Me} GC ^{5Me} T GGT GAG TTT C ^{5Me} A ^{PAL}
15	7	111496	P=S	<u>AGC</u> ^{5Me} <u>TTC</u> ^{5Me} TTT GC ^{5Me} A C ^{5Me} AT <u>GTA</u> <u>AA</u> ^{IBU}
	7	111497	P=S	<u>AGC</u> ^{5Me} <u>TTC</u> ^{5Me} TTT GC ^{5Me} A C ^{5Me} AT <u>GTA</u> <u>AA</u> ^{PAL}

¹All underlined nucleosides contain the 2'-O-
 20 Methoxyethyl group. A^{IBU} = 2'-O-Ibuprofenyl-aminohexyl-adenosine {ibuprofenyl = (4-isobutylphenyl)isopropionyl}.
 A^{PAL} = 2'-O-Palmityl-aminohexyl-Adenosine.

Table XV

25	Expected	Observed	HPLC	CE	Crude	Final
	Mass	Mass	Retention	Retention	Yield	Yield
	ISIS #	(g/mol)	(g/mol)	Time (min)	Time (min)	(ODs) (ODs)
	111494-1	6795.00	6795.43	40.87	6.27	532 136
	111495-1	6845.11	6844.08	61.80	5.87	568 163
	111496-1	7422.73	7422.98	43.79	7.54	538 282
30	111497-1	7472.84	7472.92	62.73	7.27	666 130

EXAMPLE 52

Cholesterol conjugated full 2'-O-methoxyethyl (MOE)
 phosphodiester oligonucleotide SEQ ID NO: 4 (ISIS # 16952
 unconjugated, ISIS # 16296 conjugated) TCT GAG TAG CAG

AGG AGC TC

To determine the effect of conjugating a cholesterol group at the 3'-position of a uniform 2'-MOE-phosphodiester 20mer antisense oligonucleotide both the conjugated as well
5 as the non-conjugated oligonucleotides were prepared. All of the cytosine bases were 5-methylcytosines and all ribosyl sugars were 2'-O-MOE with the exception of the 3'-terminal nucleoside having the cholesterol attached which was a 2'-hydroxycytidine. Attachment of the cholesterol group was
10 via a 6-aminohexyloxy linker at the 3' position of the conjugated oligonucleotide. The cholesterol molecule is attached to the amino group of the linker via a carbamate linkage.

The plasma concentration of the cholesterol conjugated
15 oligonucleotide (^3H , ISIS-16296) was compared to the parent oligonucleotide (^3H , ISIS-16952, Figure 2). The study was performed in male Sprague-Dawley rats using I.V. bolus administration of ^3H radiolabeled oligonucleotides. The plasma concentration was maintained at a higher level and
20 was reduced at a slower rate for the conjugated oligonucleotide.

The tissue distribution of the two radio labeled oligonucleotides was examined in Sprague-Dawley rats following I.V. bolus administration (Figure 3). Almost all
25 of the parent oligonucleotide was seen in the kidney cortex after 24 hours and only baseline amounts of oligonucleotide was seen in the other major organs tested (plasma, liver, spleen, small intestine, large intestine and mesent LN. The distribution profile for the conjugated oligonucleotide
30 showed distribution to all the organs in much higher concentrations than the parent oligonucleotide.

The percent of the dose excreted through the urine was calculated for 0-6 and 6-24 hours for the parent and conjugated oligonucleotides (Figure 4). About 38% of the

parent oligonucleotide or metabolites thereof was excreted within the first 6 hours of administration. Only about 5% of the conjugated oligonucleotide was excreted during the same time periods.

5 In a similar study, SEQ ID NO: 5 (ISIS-3082) was prepared along with 5 z for a variety of comparative pharmacokinetic studies including protein binding. The parent compound a 20mer phosphorothioate was compared with the phosphorothioate and phosphodiester 2'-propoxy analogs,
10 a chimeric analog having 2'-propoxy diester wings and a phosphorothioate deoxy center, and 5'-octadecylamine and 5'-(2'-O-hexylamino-carbonyl-oxy-cholesterol) phosphorothioate analogs. This study, in part, reported decreased excretion of the cholesterol modified oligonucleotide relative to the
15 parent phosphorothioate oligonucleotide (Crooke et al., *The Journal of Pharmacology and Experimental Therapeutics*, 1996, 277:923-937).

Example 53

Effect of conjugation on 2'-methoxyethoxy-substituted 20 phosphodiester (PO/MOE) oligonucleotides

It was observed that a PO/MOE-cholesterol conjugated oligonucleotide (16296) exhibited improved binding compared to the PO/MOE analog (16952), but is still a weak binder compared to the 2'-methoxyethoxy-substituted
25 phosphorothioate (PS/MOE) oligonucleotide (11159). The PO/MOE-ibuprofen conjugate (27700), however, not only exhibited improved binding compared to the unconjugated PO/MOE analog (16952), but also showed tighter binding than the PS analog (3067) or the PS/MOE analog (11159). These
30 results are shown below in Table XVI.

Table XVI

Oligo.	Sequence	Description	K _d (μM)
3067	TCTGAGTAGCAGAGGAGCTC	Full PS	41.2 ± 7.3
11159	TCTGAGTAGCAGAGGAGCTC	Full PS, MOE	29.3
5 16952	TCTGAGTAGCAGAGGAGCTC	Full PO, MOE	672 ± 7.21
16296	TCTGAGTAGCAGAGGAGCTC	Full PO, MOE, cholesterol conjugate	225
27700	TCTGAGTAGCAGAGGAGCTC	Full PO, MOE, ibuprofen conjugate	10.0 ± 1.30

Example 54**Conjugation of human α₁-acid glycoprotein (AAG) binding drugs
10 to oligonucleotides**

The following drug moieties were identified as drugs that bind to AAG: acenocoumarol, chlorpromazine, dipyridamole, imipramine, methadone, perphenazine, phenylbutazone, pindolol, progesterone, propanolol, RU
15 42633, RU 38486, thioridazine, ticlopidine, trifluoperazine, warfarin and phenothiazines.

Among the various phenothiazine ligands, 2-chloro-10-(2-carboxyethyl)-phenothiazine was selected as a conjugated

ligand for illustrative purposes. 2-chloro-10-(2-carboxyethyl)-phenothiazine (Melikian et al., *J. Pharm. Sci.*, **1977**, 66:228, 1977) is converted to pentafluorophenyl ester using pentafluorophenol and DCC. This compound is then condensed with 3'-O-(6-aminohexyl)-5'-O-DMT uridine and further converted to its controlled pore glass derivative. Oligonucleotides are synthesized from the controlled pore glass as described for other examples.

Example 55

10 Improved cellular uptake by conjugation of cell surface integrins with oligonucleotide

Fibrinogen-derived peptides (RGD and RGD like) are prepared for conjugation via standard peptide synthesis procedures.

15 Peptide I RIARGDFPDDRK (SEQ ID NO: 8) (an RGD peptide)

 Peptide II DELAEGGGVRGPRV (SEQ ID NO: 9)

 These peptides were synthesized in the solid phase synthesizers. At the amino terminal end, 6-hexene-20 carboxylic acid is coupled. After deprotection of the peptide, the olefinic linkage is converted into an aldehyde using OsO₄/N-methyl-morpholine oxide followed by NaIO₄ oxidation. The aldehyde containing peptide is conjugated to -O-NH₂ linked oligonucleotides. Surface plasmon resonance25 experiments indicated that these peptide conjugated oligonucleotides bind to cell surface integrins.

Example 56

Proteins and substrates to which these proteins bind

<u>Proteins</u>	<u>Substrate</u>
30 Vitamin-D binding protein	Vitamin D

Cortisol-binding globulin	Cortisol
Sex-hormone-binding protein	Sex hormones
Thyroxine-binding globulin	Thyroxine
5 and Prealbumin	

Example 57

General procedure for preparing succinimide esters of small drugs

To a solution of 1 mmol acid and 1 mmol N-hydroxy-succinimide in 4 mL of dry THF 1 mmol of dicyclohexylcarbodiimide (DCC) was added. The reaction mixture was stirred for two days at room temperature. The precipitate was filtered off and the filtrate was concentrated under reduced pressure. The residue was dissolved in 3 mL of CH_2Cl_2 , and the product purified by thick layer chromatography: Chromatotron, 2 mm plate, using a gradient from 0 to 5% of MeOH in CH_2Cl_2 . Appropriate fractions were collected, solvent removed under reduced pressure, and the product was dried under reduced pressure overnight.

20

Drug	MW acid	R _f (5% MeOH in CH_2Cl_2)		Yield (%)	MW of ester
		acid	ester		
Indomethacin	357.8	0.35	0.8	95	454.8 $\text{C}_{23}\text{H}_{19}\text{ClN}_2\text{O}_6$
Mycophenolic acid	320.3	0.25	0.9	72	417 $\text{C}_{21}\text{H}_{23}\text{NO}_8$
25 Acetyl Salicylic acid	180.2	0.1	0.6	30	277 $\text{C}_{13}\text{H}_{11}\text{NO}_6$

NHS ester of indomethacin: ^1H NMR (400 MHz, CDCl_3) δ

30 7.68 (2H, d, $J = 12.0$ Hz), 7.48 (2H, d, $J = 12.0$ Hz), 6.98 (1H, d, $J = 3.6$ Hz), 6.93 (1H, d, $J = 13.5$ Hz), 6.68 (1H, dd, $J = 13.5, 3.6$ Hz), 3.98 (2H, s), 3.87 (3H, s), 2.84 (4H,

s), 2.39 (3H, s).

NHS ester of mycophenolic acid: ^1H NMR (400 MHz, CDCl_3) δ 7.68 (1H, s), 5.34 (1H, t, $J = 10.2$ Hz), 5.21 (2H, s), 3.42 (2H, d, $J = 10.2$ Hz), 2.83 (4H, s), 2.71 (2H, t, $J = 12.0$ Hz), 2.43 (2H, t, $J = 12.0$ Hz), 2.16 (3H, s), 1.84 (3H, s).

NHS ester of acetyl salicylic acid: ES-MS: 300 $[\text{M}+\text{Na}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 8.11 (1H, dd, $J = 11.8, 2.4$ Hz), 7.66 (1H, ddd), 7.36 (1H, dd), 7.18 (1H, d, $J = 12.3$ Hz), 2.83 (4H, s), 2.30 (3H, s). ^{13}C NMR (100 MHz, CDCl_3) δ 169.0, 159.5, 151.2, 135.8, 132.1, 126.2, 124.2, 118.8, 25.4, 20.7.

The thiocresol ester of acetyl salicylic acid was prepared by stirring 180 mg (1 mmol) of acetyl salicylic acid, 124 mg (1 mmol) of *p*-thiocresol and 206 mg (1 mmol) of DCC in 4 mL of anhydrous THF at room temperature for 2 days. The precipitate was filtered off, the filtrate was concentrated under reduced pressure, the residue dissolved in 3 mL of CH_2Cl_2 and purified by preparative thick layer chromatography (Chromatotron, 2 mm plate, using a gradient from 0 to 10% MeOH in CH_2Cl_2). Obtained 256 mg (90 % yield) of white solid. R_f 0.95 in 5% MeOH in CH_2Cl_2 . $\text{C}_{16}\text{H}_{14}\text{O}_3\text{S}$. Calc: 286. ^1H NMR (400 MHz, CDCl_3) δ 8.06 (1H, dd, $J = 11.7, 2.2$ Hz), 7.59 (1H, ddd), 7.41 - 7.25 (5H, m), 7.16 (1H, d, $J = 12.0$ Hz), 2.41 (3H, s), 2.32 (3H, s).

The pentachlorophenol ester of acetyl salicylic acid was prepared by stirring 180 mg (1 mmol) of acetyl salicylic acid, 266 mg (1 mmol) pentachlorophenol and 206 mg (1 mmol) DCC in 4 mL of anhydrous THF at room temperature for 2 days. The precipitate was filtered off, the filtrate was concentrated under reduced pressure, the residue dissolved in 3 mL of CH_2Cl_2 and purified by preparative thick layer


chromatography (Chromatotron, 2 mm plate, using a gradient from 0 to 10% MeOH in CH_2Cl_2). Obtained 260 mg (61% yield) of a pale yellow solid. R_f 0.9 in 5% MeOH in CH_2Cl_2 . $\text{C}_{15}\text{H}_7\text{O}_4\text{Cl}_5$. Calc. 426. ^1H NMR (400 MHz, CDCl_3) δ 8.31 (1H, dd, $J = 11.9, 2.5$ Hz), 7.72 (1H, ddd), 7.44 (1H, ddd), 7.23 (1H, d, $J = 12.0$ Hz), 2.31 (3H, s).

Example 58

General procedure for preparing conjugated active esters of small molecules to 3'-(2'-aminopropyl)-TCT GAG TAG CAG

10 AGG AGC TC (SEQ ID NO: 4 ISIS-16952)

Solution conjugation chemistry via activated esters



Preparation on an analytical scale used a reaction volume of 500 mL, pH 9 (0.1 M $\text{NaHCO}_3/\text{Na}_2\text{CO}_3$) with the following: 10 ODs in 20 mL of H_2O of **SEQ ID NO: 4**-3'- NH_2 in triethylammonium form (TEA); 100 mL of 0.5 M buffer and 255 mL DMSO. To this solution, 125 mL of 0.05 M solution of activated ester in DMSO was added. The mixture was vortexed and the resultant heterogeneous mixture was kept at room temperature. The aliquots were analyzed by RP- C_{18} HPLC after

2, 4, 6 and 18 h. It was shown that the reaction was completed at 6 hours. The conjugates had higher retention times than the oligonucleotide.

	Modification	Retention time (min)	ES-MS		Yield% (HPLC)	ISIS #
			Calculated	Found		
5	SEQ ID NO: 4-3'-NH ₂	16				
	SEQ ID NO: 4-indomethacin	22	8046.8	8048.17	85	124546
	SEQ ID NO: 4-mycophenolic acid	18.5	8009	8009.5	80	124545
10	SEQ ID NO: 4-acetylsalicylic acid	17.5	7869	7866	60	30785
	SEQ ID NO: 4-salicylic acid	18	7826	7824.8		126785
	SEQ ID NO: 4-NH-Ac	16.5				

15 **HPLC conditions:** RP-C₁₈ Delta Pak column, 3.9 mm x 300 mm, 15 m, 1.5 mL/min. A: 0.1 M NH₄OAc; B: 80% MeCN Gradient from 0 to 60% B in 30 min.

Example 59

Conjugation of acetylsalicylic acid

20 Various activated esters of acetylsalicylic acid were used to conjugate aspirin. A better yield was obtained using pentachlorophenol ester which gave the aspirin conjugate in 60% yield. For some activated esters, that were examined in this reaction sequence, only acetylation of
25 the amino group of the oligonucleotide was observed.

Activated ester	Yield of SEQ ID NO: 4-NH-Ac	Yield of SEQ ID NO: 4-aspirin
NHS	100%	
p-Thiocresol	100%	
Pentachlorophenol	40%	60%

5

Example 60**Stability of SEQ ID NO: 4-acetylsalicylic acid conjugate**

The stability of SEQ ID NO: 4-aspirin conjugate was
10 examined in various solution. The stability of the
conjugate in various conditions was estimated by the
analysis of conjugate solutions by RP-C₁₈ HPLC at different
incubation times.

Buffer	pH	T _{1/2}
Water	8.2	10 days
0.05 M Triethylammonium acetate	7.0	4 days
1 M NH ₄ OAc	6.8	6 h
0.1 M NH ₄ OAc	6.0	4 days
0.1 M NH ₄ OAc	6.5	48 h
0.1 M NH ₄ OAc	7.0	24 h
0.1 M NH ₄ OAc	7.5	12 h

15

20

Example 61**25 Preparative Scale synthesis of conjugates**

In Preparative scale, a heterogeneous solution of 100
ODs of SEQ ID NO: 4-3'-NH₂ in TEA form, 100 mL 0.5 M buffer
and 400 mL 0.1 M solution of activated ester in DMSO were
mixed and vortexed for 6h at room temperature. Excess
30 activated ester was removed by gel filtration through
Sephadex G-25. The conjugate was purified and desalted by
RP-C₁₈ HPLC. Yield: from 30 to 50 ODs of the conjugate.


Example 62**ISIS SEQ ID NO: 4-salicylic acid conjugate**

The title conjugate was obtained by hydrolysis of SEQ ID NO: 4-aspirin conjugate in 0.1 M NH_4OAc , pH 7.5, 18 h.

Example 63

Warfarin conjugation

- 5 I. Synthesis of C6 aminooxy linker and its phosphoramidite was by the following scheme:



Compound 2

1,6-Hexanediol, 8.5 g (72 mmol) was dried first by coevaporation with anhydrous pyridine (2 x 50 mL), then overnight under high vacuum. The residue was dissolved in a mixture of 60 mL dry 1,4-dioxane and 10 mL of dry Py and 8.0 g (23.6 mmol) DMT-Cl was added in 4 portions. The reaction mixture was stirred under Ar at RT for 2 days. The solution was concentrated under reduced pressure, the residue was

dissolved in 300 mL of CH_2Cl_2 , washed with 5% aq NaHCO_3 , then brine. The organic phase was concentrated under reduced pressure. A yellow oil (9 g, 21.4 mmol) was obtained after purification by column chromatography using a gradient of 5 MeOH, from 0 to 10%, in 0.2% $\text{Py}/\text{CH}_2\text{Cl}_2$. ^1H NMR (400 MHz, CDCl_3) δ 7.46-7.43 (2H, m), 7.35-7.26 (5H, m), 7.20-7.17 (2H, m), 6.85-6.80 (4H, m), 3.80 (3H, s), 3.79 (3H, m), 3.65 (2H, t, $J = 7.2$ Hz), 3.1-3.0 (2H, m), 1.63-1.55 (4H, m), 1.42-1.34 (4H, m). ^{13}C NMR (100 MHz, CDCl_3) δ 158.8, 158.5, 145.6, 139.7, 136.9, 130.2, 129.4, 128.1, 128.0, 127.9, 113.4, 113.2, 85.8, 63.6, 63.2, 55.4, 32.9, 30.3, 26.3, 25.8.

Compound 3

Monosubstituted hexanediol (**Compound 2**), (1.68 g, 4 mmol), (0.7 g, 4.3 mmol) of *N*-hydroxyphthalimide and (1.4 g, 4.2 mmol) of PPh_3 was dissolved in anhydrous dioxane. To this solution 0.7 mL (3.6 mmol) of diethylazodicarboxylate (DEAD) was added dropwise, and the reaction mixture stirred overnight at room temperature. TLC in 5% MeOH/ CH_2Cl_2 , 1% Et_3N indicated the completion of reaction (a new spot with R_f 0.9, compared to R_f 0.6 of **Compound 2**). Reaction mixture was concentrated under reduced pressure. The residue was dissolved in hexane/ethylacetate, 4/1, by an addition of CH_2Cl_2 . The product was purified by preparative column chromatography, eluted with 1% Et_3N , hexane/ethylacetate, 3/2, to give after drying under reduced pressure 2 g (89%) of colorless glass. ^1H NMR (400 MHz, CDCl_3) δ 7.83-7.81 (2H, m), 7.74-7.72 (2H, m), 7.42 (2H, d, $J = 8.4$ Hz), 7.35 - 7.25 (6H, m), 7.20 - 7.18 (1H, m), 6.81 (4H, d, $J = 8.8$ Hz), 4.17 (2H, t, $J = 6.4$ Hz), 3.77 (6H, s), 3.04 (2H, t, $J = 6.4$ Hz), 1.79 - 1.75 (2H, m), 1.64 - 1.57 (2H, m), 1.44 - 1.42 (4H, m). ^{13}C NMR (100 MHz, CDCl_3) δ 164.0, 158.2, 145.6, 136.8, 134.4, 130.2, 129.7, 128.5, 128.0, 126.7, 123.6, 113.2, 85.8, 78.3, 63.4, 55.6, 30.6, 28.2, 26.1, 25.8.

Compound 4

Compound 3, 1.1 g (1.95 mmol) was dissolved in 20 mL of $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 9:1, the dichloroacetic acid was added until stable red color appeared and then kept for at room temperature for an additional 45 min. The reaction mixture was quenched with 50 mL of 5% aq. NaHCO_3 and diluted with 50 mL CH_2Cl_2 . The organic phase was washed with 50 mL of brine and concentrated under reduced pressure. The product was purified by preparative thick layer chromatography (Chromatotron, plate 2 mm). Appropriate fractions were collected and dried under reduced pressure to give 0.36 g (70%) of white solid. R_f 0.6 in 5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$, 1% Et_3N . MS (positive MeOH): 264.2 $[\text{M} + \text{H}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 7.84-7.82 (2H, m), 7.75-7.73 (2H, m), 4.21 (2H, t, $J = 6.6$ Hz), 3.69-3.66 (2H, m), 1.81-1.79 (2H, m), 1.63-1.53 (4H, m), 1.47-1.44 (2H, m). ^{13}C NMR (100 MHz, CDCl_3) δ 163.9, 134.6, 129.2, 123.7, 78.5, 62.9, 32.7, 28.3, 25.5.

Compound 5


To a solution of **Compound 4**, 300 mg (1.14 mmol) in 4 mL of anhydrous MeCN under Ar, 543 mL (1.75 mmol, 1.5 eq) of phosphoramidite reagent (2-cyanoethyl tetraisopropylphosphorodiamidite) and 2.28 mL (1.03 mmol, 0.9 eq.) of 0.45 M solution of tetrazole in MeCN were added. A precipitate appeared in a few seconds. After 1h, 300 mL of Et_3N was added to the reaction mixture followed by quenching with 30 mL of 5% aq. NaHCO_3 . The product was rapidly extracted with 50 mL CH_2Cl_2 , the organic phase was washed with 30 mL brine, then 30 mL of water, concentrated under reduced pressure. The product was purified by preparative thick layer chromatography (Chromatotron, plate 2 mm) in the gradient from 5 to 50% ethyl acetate in hexane with 1% Et_3N . Appropriate fractions were collected and dried under reduced pressure to give 188 mg (35%) of colorless oil. R_f 0.4 in

hexane/ethylacetate/Et₃N, 70/30/5. ³¹P NMR (162 MHz) δ 148.2, 148.0.

Example 64

Synthesis of Warfarin phosphoramidite

5



Compound 7

To 1 mmol of **Compound 3** in 6 mL of anhydrous CH_2Cl_2 at -10°C was added 1.1 eq. of methylhydrazine which formed a white precipitate. After 1h an additional 5 mL of cold CH_2Cl_2 was added, and the reaction mixture was rapidly filtered. The precipitate was washed with 5 mL of cold CH_2Cl_2 . Warfarin (308 mg, 1 mmol) was added to the filtrate and the solution was allowed to stand at room temperature for 1h. TLC showed a new fluorescent spot with R_f 0.8 in 2% MeOH, 0.1% Et_3N in CH_2Cl_2 . The reaction mixture was concentrated under reduced pressure and the crude material was purified by preparative thick layer chromatography (Chromatotron, plate 2 mm) using a gradient from 0 to 10% MeOH in CH_2Cl_2 , 1% Et_3N . Obtained 0.36 g (50%) of colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.89 (2H, d, J = 8.4 Hz), 7.48–7.41 (4H, m), 7.32 – 7.24 (6H, m), 7.20 – 7.06 (6H, m), 6.81 (4H, d, J = 8.8 Hz), 4.87 – 4.84 (1H, m), 3.92 (2H, t, J = 7.2 Hz), 3.78 (6H, s), 3.28 – 3.24 (1H, m), 3.14 – 3.09

(1H, m), 2.99 (2H, t, $J = 6.4$ Hz), 1.85 (3H, s), 1.58 - 1.46 (4H, m), 1.27 - 1.19 (4H, m).

Compound 8

Pivaloyl chloride, 84 mL (0.7 mmol), was added to a solution of **Compound 7**, 400 mg (0.55 mmol) and 20 mg DMAP in 10 mL of anhydrous pyridine. A precipitate appeared immediately after addition, and the heterogeneous reaction mixture was stirred for 2 h at room temperature. The TLC in 0.5% MeOH, 0.1% Et₃N in CH₂Cl₂ showed complete conversion of starting material to more hydrophobic one with R_f 0.45. The reaction was quenched with 50 mL of 5% aq. NaHCO₃ and the product was extracted with 2 x 50 mL of CH₂Cl₂. The organic phase was washed with 50 mL brine, then concentrated under reduced pressure and dried to give 440 mg of an oil. The oil was detritylated without further purification.

Detritylation

The oil, 440 mg, was dissolved in 20 mL CH₂Cl₂/MeOH, 9/1, and dichloroacetic acid was added until the color of the solution stabilized. TLC: a new more polar spot appeared with R_f 0.4 in 5% MeOH/CH₂Cl₂. After addition of 100 mL of 5% aq. NaHCO₃, the product was extracted with 50 mL of CH₂Cl₂. Organic phase was washed with 40 mL of brine and concentrated under reduced pressure. The product was obtained after purification by preparative thick layer chromatography (Chromatotron, plate 2 mm) using a gradient from 0 to 5% MeOH in CH₂Cl₂. Yield 200 mg (71%) of an oil. NMR spectrum shows 2 signal's sets in ratio 80/20. MS (positive MeOH): 508 [M + H]⁺. ¹H NMR (400 MHz, CDCl₃) δ 7.49-7.40 (4H, m), 7.32 - 7.19 (5H, m), 4.58 (1H, t, $J = 7.6$ Hz), 3.91 (2H, t, $J = 6.8$ Hz), 3.58 (2H, t, $J = 6.8$ Hz), 3.20 (2H, bs), 1.82 (3H, s), 1.54 (9H, s), 1.48 - 1.43 (4H, m), 1.23 - 1.21 (2H, m). ¹³C NMR (100 MHz, CDCl₃) δ 175.1, 160.9, 154.8, 152.7, 140.5, 132.0, 128.4, 126.9, 124.5, 122.9,

122.1, 116.95, 116.4, 73.5, 63.0, 39.9, 38.9, 37.3, 32.9, 29.2, 27.6, 25.9, 25.8, 25.7, 15.2.

Compound 9

Compound 8, 0.18 g (0.36 mmol), was dissolved in 5 mL of anhydrous MeCN. To this solution 169 mL (0.53 mmol, 1.5 eq.) of phosphoramidite reagent and 0.71 mL (0.32 mmol, 0.9 eq.) of 0.45 M 1H-tetrazole solution in MeCN were added. The reaction mixture was stirred under Ar for 3 h at room temperature. TLC in hexane/ethylacetate/Et₃N, 70/30/5, 10 revealed a new spot with R_f 0.6, as well as fluorescent spot with R_f 0.5 which resulted from a loss of pivaloyl protecting group. 200 mL of Et₃N followed by 30 mL of 5% aq. NaHCO₃ solution was added to the reaction mixture. The product was rapidly extracted with 50 mL CH₂Cl₂, organic phase was washed 15 with 30 mL of brine, 20 mL of water and concentrated under reduced pressure. After purification by preparative thick layer chromatography (Chromatotron, plate 2 mm) using a gradient from 0 to 10% MeOH in CH₂Cl₂, 1% Et₃N, 186 mg (0.26 mmol, 72%) of a colorless oil was obtained. ³¹P NMR (161 20 MHz, MeCN) δ 148.2, 148.0. The purity was greater than 98%.

Example 65

Warfarin containing controlled-pore glass CPG 10

Tritylation

2'-O-(ethyloxypthalimido)5-methyluridine, 223 mg (0.5 25 mmol), was dried by coevaporation with 50 mL of anhydrous MeCN, then overnight under reduced pressure. The dried compound was dissolved in 5 mL of anhydrous pyridine and 203 mg (0.6 mmol) of DMTCl was added. After 1 h at room temperature, TLC showed the reaction was incomplete. 30 Additional DMTCl, 40 mg (0.1 mmol), was added. After 30 min 50 mL of 5% aq. NaHCO₃ was added and the product was extracted with 2 x 60 mL CH₂Cl₂. The organic layer was dried

under reduced pressure. The product was purified by preparative thick layer chromatography (Chromatotron, plate 2 mm) using a gradient from 0 to 10% MeOH in CH_2Cl_2 , 3% Et_3N . The appropriate fractions were collected and dried under reduced pressure to give 100 mg (26%) of a pale yellow oil. FW 749. ^1H NMR (400 MHz, CDCl_3) δ 9.65 (1H, s), 7.89-7.76 (5H, m), 7.48 - 7.27 (9H, m), 6.88 (4H, d, J = 8.4 Hz), 6.00 (1H, bs), 4.61 - 4.42 (3H, m), 4.36 - 4.05 (4H, m), 3.83 (6H, s), 3.59 - 3.54 (2H, m), 2.80 - 2.78 (1H, m), 1.39 (3H, s).

10 **Succinic ester**

To a solution of 170 mg (0.23 mmol) of the DMT-nucleoside in 1 mL anhydrous 1,2-dichloroethane, 15 mg (0.125 mmol, 0.5 eq.) of DMAP, 35 mL (0.25 mmol, 1.1 eq.) Et_3N , and 34 mg (0.34 mmol, 1.5 eq) succinic anhydride were added. The reaction was heated at 50°C for 30 min under anhydrous conditions. 1,2-Dichloroethane, 15 mL, was added and the solution was washed with 2 x 10 mL of ice-cold 10% citric acid and 2 x 10 mL of water. The organic layer was concentrated and dried under high vacuum to give 149 mg of a yellow foam. This foam was dissolved in 2 mL of CH_2Cl_2 , and precipitated in 50 mL of hexane/ether, 1/1. v/v. White powder was obtained: 87 mg (45%). FW 849. $\text{C}_{45}\text{H}_{43}\text{N}_3\text{O}_{14}$. R_f 0.3 in 5% MeOH, 0.1% Et_3N in CH_2Cl_2 . ^1H NMR (400 MHz, CDCl_3) δ 9.63 (1H, s), 7.78-7.67 (5H, m), 7.39 - 7.22 (9H, m), 6.84 - 6.81 (4H, m), 5.95 (1H, d, J = 3.2 Hz), 5.33 (1H, t, J = 6.0 Hz), 4.48 - 4.38 (2H, m), 4.27 (1H, t, J = 4.8 Hz), 4.18 - 4.10 (2H, m), 3.96 - 3.91 (1H, m), 3.78 (6H, s), 3.50 (1H, d, J = 10.6 Hz), 3.26 (1H, dd, J = 10.8, 2.0 Hz), 2.83 - 2.48 (4H, m), 1.37 (3H, s).

30 **Loading**

The nucleoside succinate, 87 mg (0.1 mmol), and 12.2 mg (0.1 mmol) of DMAP in 1 mL of anhydrous MeCN was added to a solution of 38 mg (0.12 mmol) of 2,2'-dithiobis(5-

nitropyridine) in 1 mL of MeCN/CH₂Cl₂, 0.75/0.25, v/v. To this mixture, a solution of 26 mg (0.1 mmol) of triphenylphosphine in 0.5 mL of MeCN was added to give a red solution. After a brief agitation 0.5 g of acid-washed LCAA-CPG was added. The suspension was further agitated for 1h. The resin was successively washed with 20 mL portions of MeCN, CH₂Cl₂ and diethyl ether followed by drying. The residual amino groups were capped by treatment with 2 mL of CapA and CapB solutions for 2h. The resin was washed with MeCN and dried under reduced pressure. The nucleoside loading was determined by trityl test: 6.5 mg of resin in 30 mL of 3% dichloroacetic acid in CH₂Cl₂. Absorbance of DMT⁺ was measured at 504 nm, $e = 76 \text{ mL} \times \text{mmol}^{-1} \times \text{cm}^{-1}$. Obtained loading 72.8 mmol/g.

15 CPG 11

10 mg of CPG-10 was treated with 1 mL of 0.5 M hydrazine acetate in pyridine for 1 h at room temperature, then washed successively with 1 mL of pyridine, methanol, MeCN, and dried. A solution of warfarin (1mL, 1M) in DMF was added to resin, the mixture was shaken for 24 h at room temperature, washed with 3 x 1 mL of DMF, 2 x 1 mL MeCN and dried.

Analysis

detritylation with 3% CF₃COOH/ CH₂Cl₂;
25 cleavage with 0.5 mL of concentrated aqueous ammonia, 1h, room temperature;
RP-HPLC, Delta Pak C₁₈, 3.9 x 300 mm, flow 1.5 mL/min
A: 0.1 M NH₄OAc; B: 80% MeCN, Gradient from 0 to 60% B in 30 min.

30 2'-group	Retention time	FW for 5',3'-
OH-T		
Aminooxyethyl	12.1 min.	317

Phthalimidooxyethyl	20.1 min.	447
Warfarinoaminooxyethyl	20.8 min.	607
Acetoaminooxyethyl	16.3 min.	357

After RP-HPLC analysis, warfarin adduct was obtained
5 (80% yield) with 10% of the acetone adduct. MS (positive
MeOH): 608.2 [M+H]⁺. Reference acetone adduct was prepared
by reaction of aminooxyethyl nucleoside after cleavage of
phthalimide group by hydrazine acetate with acetone.

CPG 12

- 10 CPG 11, 30 mg, was treated (syringe technique) with 1 M
Pivaloyl chloride, 0.05 M DMAP in pyridine for 30 min. Then
the resin was washed with 5 mL of pyridine, 10 mL MeCN,
dried.

Example 66

- 15 Post-oligomerization conjugation of warfarin to 5'-terminus
of an oligonucleotide (T₁₂) (SEQ ID NO: 10)

Phosphoramidite 5 was used for incorporation of an
aminoxy linker at the 5'-position of a T₁₂ oligonucleotide.
Coupling conditions: 0.1 M amidite solution, 1.9 mL of
20 amidite per couple, 17 min couple time, CSO oxidation, 4
min. 3 mg of resin was treated with conc. aqueous ammonia
for 1 h at room temperature. ES-MS: expected 3783.8; found
3783.75 (for phthalimide deprotected oligo).

Warfarin coupling

- 25 Resin with aminoxy linker, 40 mg, was treated (syringe
technique) with 1 mL of 0.5 M hydrazine acetate solution for
30 min. The resin was washed with 5 mL pyridine, 5 mL of
MeOH, 10 mL of MeCN and dried. The resin was placed in an
ependorf tube, and 1 mL of a 0.15 M warfarin solution in
30 MeCN/DMF, 2/1, v/v was added. The mixture was shaken for 24
h, the resin washed with 1 mL DMF, 3 x 1 mL MeCN and dried.
Cleavage in conc. aq. ammonia 1h at room temperature and 8 h

at 55°C gave similar RP-HPLC profiles and ES-MS spectra: Expected 4073, Found 4073. Hence the warfarin-conjugate showed stability under standard cleavage/deprotection conditions. After HPLC purification the conjugation efficiency was >95%.

Example 67

Incorporation of warfarin phosphoramidite at 5'-end of oligonucleotide

Full diester thymidine 12mer (T_{12} , $P=O$), 1.5 mmol scale, 0.1 M amidite solution, 380 mL of solution per couple, coupling time 14 min, CSO oxidation. After cleavage in conc. aq. ammonia, ES-MS: Expected 4073, Found 4074. HPLC: RP C_{18} , A: 0.1 M NH_4OAc ; B: 80% MeCN, Gradient from 0 to 60% B in 60 min.

5'-Warfarin-3082 was prepared from the following conditions:
1 mmol scale, 14 min coupling, cleavage and deprotection conc. aq. ammonia, 8 h, 55°C. After HPLC, synthesis was not bad. ES-MS: expected 6469.7; found 6469.5. After purification 16 ODs were obtained.

Example 68

Warfarin conjugates

Post-oligomerization conjugation

2 mmol synthesis of SEQ ID NO. 5 starting with CPG-10, bearing 2'-phthalimidooxyethyl-T, was performed through standard CSO protocol. The resin was treated with 1 mL of 0.5 M hydrazine acetate solution for 1 h, then washed with 5 mL Py, 5 mL MeOH, 10 mL MeCN, dried under reduced pressure for 15 min. 1 mL of 1 M solution of warfarin in DMF, was added, and mixture was shaken overnight. The solution was decanted, the resin washed 2 x 1 mL DMF, 3 x 1 mL MeCN, and

dried. Cleavage and deprotection was effected using conc. aq. ammonia for 8 h at 55°C. HPLC and ES-MS showed two major products: 60% of warfarin adduct (calc. 6651 (DMT-on); found 6652), and 40% of the acetone adduct (6401). Obtained 5 30 ODs after HPLC purification and detritylation. ES-MS: expected 6349; found 6349.

SEQ ID NO. 5-3'-Warfarin was synthesized in similar way. Obtained 10 ODs of purified oligonucleotide (should be able to increase this yield with optimization of HPLC 10 conditions). ES-MS: expected 8011, found 8013.

Synthesis using CPG 12

The oligonucleotide was synthesized using 14 mg of resin (CPG 12), 0.15 M amidite solutions and CSO oxidation. Cleavage and deprotection: 8 h at 55°C in conc. aq. ammonia. 15 gave SEQ ID NO: 4 sequence. After HPLC purification in trityl-on and purification in trityl-off form, 12 ODs of oligonucleotide were obtained. ES-MS: expected 8011, found 8014.

Example 69

20 Effect of Chemistry on α_2 -Macroglobulin Binding

The binding affinity of different chemistries was evaluated with α_2 -macroglobin as the target protein using the techniques that are described in Example 42.

25	<u>K_d (μM)</u>	<u>Description (analogs of SEQ ID No. 4)</u>
	0.43	PS deoxy
	0.20	PS 2'-O-propyl
	5.7	PO 2'-O-methoxyethyl/chlosterol
	0.75	PS 2'-O-methoxyethyl
30	13.8	PO 2'-O-methoxyethyl/ibuprofen
	1.9	PO deoxy/palmityl
	no binding	PO 2'-deoxy

no binding PO 2'-O-propyl

As can be seen from the tables even with the PO backbone the oligomers exhibit good binding affinity when they have ligands such as cholesterol, ibuprofen and 5 palmitic acid. Among these ligands palmitic acid shows favorable binding to α_2 -macroglobin while ibuprofen shows favorable to serum albumin.

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204